

**YWHAG Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AW5242****Specification**

---

**YWHAG Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">P61981</a>
Other Accession	<a href="#">Q6NRY9</a> , <a href="#">Q6PCG0</a> , <a href="#">Q6PC29</a> , <a href="#">P61983</a> , <a href="#">P61982</a> , <a href="#">Q5F3W6</a> , <a href="#">P68252</a>
Reactivity	Human, Mouse
Predicted	Bovine, Chicken, Rat, Zebrafish, Xenopus
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=28,M=28,Rat=28 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

**YWHAG Antibody (N-term) - Additional Information****Gene ID** 7532**Antigen Region**  
63-92**Other Names**

YWHAG; 14-3-3 protein gamma; Protein kinase C inhibitor protein 1; 14-3-3 protein gamma, N-terminally processed

**Dilution**WB~~1:1000  
IHC-P~~1:10~50  
FC~~1:10~50**Target/Specificity**

This YWHAG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-92 amino acids from the N-terminal region of human YWHAG.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

YWHAG Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## YWHAG Antibody (N-term) - Protein Information

**Name** YWHAG

### Function

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/15696159" target="\_blank">15696159</a>, PubMed:<a href="http://www.uniprot.org/citations/16511572" target="\_blank">16511572</a>, PubMed:<a href="http://www.uniprot.org/citations/36732624" target="\_blank">36732624</a>). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:<a href="http://www.uniprot.org/citations/15696159" target="\_blank">15696159</a>, PubMed:<a href="http://www.uniprot.org/citations/16511572" target="\_blank">16511572</a>, PubMed:<a href="http://www.uniprot.org/citations/36732624" target="\_blank">36732624</a>). Binding generally results in the modulation of the activity of the binding partner (PubMed:<a href="http://www.uniprot.org/citations/16511572" target="\_blank">16511572</a>). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed:<a href="http://www.uniprot.org/citations/36732624" target="\_blank">36732624</a>).

### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P68252}.

### Tissue Location


Highly expressed in brain, skeletal muscle, and heart.


## YWHAG Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## YWHAG Antibody (N-term) - Images






Western blot analysis of lysates from HeLa, A431 cell line, mouse brain tissue lysate (from left to right), using YWHAG Antibody (N-term) (Cat. #AW5242). AW5242 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody.



Formalin-fixed and paraffin-embedded human brain tissue reacted with YWHAG Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



YWHAG Antibody (N-term) (Cat. #AW5242) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **YWHAG Antibody (N-term) - Background**

YWHAG belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 100% identical to the rat ortholog. It is induced by growth factors in human vascular smooth muscle cells, and is also highly expressed in skeletal and heart muscles, suggesting an important role for this protein in muscle tissue. It has been shown to interact with RAF1 and protein kinase C, proteins involved in various signal transduction pathways.

#### **YWHAG Antibody (N-term) - References**

Jagemann, L.R., et al., J. Biol. Chem. 283 (25), 17450-17462 (2008)