10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HSP90B1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5268

Specification

HSP90B1 Antibody (Center) - Product Information

Application	**WB, IHC-P, FC,E**
Primary Accession	P14625
Other Accession	Q4R520, NP_003290.1
Reactivity	**Human, Rat**
Predicted	**Monkey**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**H=92;M=92;Rat=93,74 KDa**
Isotype	**Rabbit IgG**
Antigen Source	**HUMAN**

HSP90B1 Antibody (Center) - Additional Information

Gene ID 7184

Antigen Region
460-487

Other Names
HSP90B1; GRP94; TRA1; Endoplasmin; 94 kDa glucose-regulated protein; Heat shock protein 90 kDa beta member 1; Tumor rejection antigen 1; gp96 homolog

Dilution
WB~~ 1:1000
IHC-P~~1:10~50
FC~~1:10~50

Target/Specificity
This HSP90B1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 460-487 amino acids from the Central region of human HSP90B1.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
HSP90B1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HSP90B1 Antibody (Center) - Protein Information

Name HSP90B1 (HGNC:12028)

Synonyms GRP94, TRA1

Function
Molecular chaperone that functions in the processing and transport of secreted proteins (By similarity). When associated with CNPY3, required for proper folding of Toll-like receptors (By similarity). Functions in endoplasmic reticulum associated degradation (ERAD) (PubMed:18264092). Has ATPase activity (By similarity). May participate in the unfolding of cytosolic leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1 to facilitate their translocation into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) and secretion; the translocation process is mediated by the cargo receptor TMED10 (PubMed:32272059).


Cellular Location
Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen {ECO:0000250|UniProtKB:P41148}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

HSP90B1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HSP90B1 Antibody (Center) - Images




10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Western blot analysis of lysates from mouse C2C12,MCF-7,Hela cell line (from left to right), using HSP90B1 Antibody (Center)(Cat. #AW5268). AW5268 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.



HSP90B1 Antibody (Center) (Cat. #AW5268)immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of HSP90B1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



HSP90B1 Antibody (Center) (Cat. #AW5268) flow cytometric analysis of A549 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HSP90B1 Antibody (Center) - Background

HSP90 proteins are highly conserved molecular chaperones that have key roles in signal transduction, protein folding, protein degradation, and morphologic evolution. HSP90 proteins normally associate with other cochaperones and play important roles in folding newly synthesized proteins or stabilizing and refolding denatured proteins after stress. HSP90B1 is an endoplasmic reticulum HSP90 protein. Other HSP90 proteins are found in cytosol (see HSP90AA1; MIM 140571) and mitochondria (TRAP1; MIM 606219) (Chen et al., 2005 [PubMed 16269234]).

HSP90B1 Antibody (Center) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)

Suzuki, S., et al. Biochem. Biophys. Res. Commun. 398(3):525-531(2010)
Bloor, S., et al. Proc. Natl. Acad. Sci. U.S.A. 107(15):6970-6975(2010)
Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Wang, X., et al. Clin. Dev. Immunol. 2010, 212537 (2010) :