

SOX2 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5343

Specification

SOX2 Antibody (C-term) - Product Information

Application WB,E
Primary Accession P48431

Other Accession
Reactivity
Predicted
Host
Clonality
P48430, P54231
Human, Mouse
Chicken, Sheep
Rabbit
Polyclonal

Calculated MW H=34;M=34 KDa Isotype Rabbit IgG Antigen Source HUMAN

SOX2 Antibody (C-term) - Additional Information

Gene ID 6657

Antigen Region

282-316

Other Names

Transcription factor SOX-2, SOX2

Dilution

WB~~1:1000

Target/Specificity

This SOX2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 282-316 amino acids from the C-terminal region of human SOX2.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SOX2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SOX2 Antibody (C-term) - Protein Information



Name SOX2

Function

Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206 (By similarity). Binds to the proximal enhancer region of NANOG (By similarity). Critical for early embryogenesis and for embryonic stem cell pluripotency (PubMed:18035408). Downstream SRRT target that mediates the promotion of neural stem cell self-renewal (By similarity). Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity). May function as a switch in neuronal development (By similarity).

Cellular Location

Nucleus speckle {ECO:0000250|UniProtKB:Q05738}. Nucleus {ECO:0000250|UniProtKB:Q05738}. Nucleus {ECO:0000250|UniProtKB:Q05738}. Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (By similarity). Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH) (By similarity) Colocalizes with SOX6 in speckles. Colocalizes with CAML in the nucleus (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (By similarity) {ECO:0000250|UniProtKB:Q05066, ECO:0000250|UniProtKB:Q05738}

SOX2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
 SOX2 Antibody (C-term) Images





Western blot analysis of lysates from F9,NCCIT cell line (from left to right), using SOX2 Antibody (C-term)(Cat. #AW5343). AW5343 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody.Lysates at 20ug per lane.

SOX2 Antibody (C-term) - Background

Transcription factor that forms a trimeric complex with OCT4 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206 (By similarity). Critical for early embryogenesis and for embryonic stem cell pluripotency. May function as a switch in neuronal development. Downstream SRRT target that mediates the promotion of neural stem cell self-renewal (By similarity). Keeps neural cells undifferentiated by counteracting the activity of proneural proteins and suppresses neuronal differentiation (By similarity).

SOX2 Antibody (C-term) - References

Stevanovic M.,et al.Mamm. Genome 5:640-642(1994).
Sadler L.A.,et al.Submitted (DEC-1992) to the EMBL/GenBank/DDBJ databases.
Fantes J.,et al.Nat. Genet. 33:461-463(2003).
Takahashi K.,et al.Cell 131:861-872(2007).
Rigbolt K.T.,et al.Sci. Signal. 4:RS3-RS3(2011).