10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CHCHD3 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5353

Specification

CHCHD3 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q9NX63
Reactivity	**Human**
Host	**Rabbit**
Clonality	**polyclonal**
Calculated MW	**H=26 KDa**
Isotype	**Rabbit IgG**
Antigen Source	**HUMAN**

CHCHD3 Antibody (Center) - Additional Information

Gene ID 54927

Antigen Region
83-116

Other Names
Coiled-coil-helix-coiled-coil-helix domain-containing protein 3, mitochondrial, CHCHD3, MINOS3

Dilution
WB~~1:1000

Target/Specificity
This CHCHD3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 83-116 amino acids from the Central region of human CHCHD3.

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CHCHD3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CHCHD3 Antibody (Center) - Protein Information

Name CHCHD3

Synonyms MIC19, MINOS3

Function
Component of the MICOS complex, a large protein complex of the mitochondrial inner membrane that plays crucial roles in the maintenance of crista junctions, inner membrane architecture, and formation of contact sites to the outer membrane. Has also been shown to function as a transcription factor which binds to the BAG1 promoter and represses BAG1 transcription. Plays an important role in the maintenance of the MICOS complex stability and the mitochondrial cristae morphology (PubMed:25781180).

Cellular Location
Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q9CRB9}; Lipid-anchor; Intermembrane side {ECO:0000250|UniProtKB:Q9CRB9}. Cytoplasm. Nucleus Mitochondrion


Tissue Location
Detected at low levels in brain, placenta, lung, liver, kidney and pancreas with increased levels in heart and skeletal muscle. Higher expression in primary lung cancers than in normal lung tissue.

CHCHD3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CHCHD3 Antibody (Center) - Images



All lanes : Anti-CHCHD3 Antibody (Center)(AW5353) at 1/1000 dilution Lane 1: 293T whole cell lysates Lane 2: A431 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 28 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.

CHCHD3 Antibody (Center) - Background

Required for maintenance of mitochondrial crista integrity and mitochondrial function. May act as a scaffolding protein that stabilizes protein complexes involved in crista architecture and protein import. Has also been shown to function as a transcription factor which binds to the BAG1 promoter and represses BAG1 transcription.

CHCHD3 Antibody (Center) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Daub H.,et al.Mol. Cell 31:438-448(2008).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).
Mayya V.,et al.Sci. Signal. 2:RA46-RA46(2009).
Choudhary C.,et al.Science 325:834-840(2009).