

**ALDH1A1 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AW5387****Specification**

---

**ALDH1A1 Antibody - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">P00352</a>
Other Accession	<a href="#">NP_000680.2</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	H=55 KDa
Isotype	Mouse IgG1
Antigen Source	HUMAN

**ALDH1A1 Antibody - Additional Information****Gene ID** 216**Antigen Region**  
7-306**Other Names**

Retinal dehydrogenase 1, RALDH 1, RaIDH1, ALDH-E1, ALHDII, Aldehyde dehydrogenase family 1 member A1, Aldehyde dehydrogenase, cytosolic, ALDH1A1, ALDC, ALDH1, PUMB1

**Dilution**

WB~~1:1000  
FC~~1:25

**Target/Specificity**

This ALDH1A1 monoclonal antibody is generated from mouse immunized with ALDH1A1 recombinant protein.

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ALDH1A1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ALDH1A1 Antibody - Protein Information**

**Name** ALDH1A1 ([HGNC:402](#))

### Function

Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide range of aldehydes to their corresponding carboxylic acid (PubMed:[19296407](http://www.uniprot.org/citations/19296407), PubMed:[12941160](http://www.uniprot.org/citations/12941160), PubMed:[15623782](http://www.uniprot.org/citations/15623782), PubMed:[17175089](http://www.uniprot.org/citations/17175089), PubMed:[26373694](http://www.uniprot.org/citations/26373694), PubMed:[25450233](http://www.uniprot.org/citations/25450233)). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (By similarity). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (By similarity). Also oxidizes aldehydes resulting from lipid peroxidation like (E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:[19296407](http://www.uniprot.org/citations/19296407), PubMed:[12941160](http://www.uniprot.org/citations/12941160), PubMed:[15623782](http://www.uniprot.org/citations/15623782)). Functions also downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3-deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycation agent that may react with lysine and arginine side-chains of proteins (PubMed:[17175089](http://www.uniprot.org/citations/17175089)). Has also an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).

### Cellular Location

Cytoplasm, cytosol. Cell projection, axon {ECO:0000250|UniProtKB:P24549}

### Tissue Location


Expressed by erythrocytes (at protein level).


## ALDH1A1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## ALDH1A1 Antibody - Images





All lanes : Anti-ALDH1A1 Antibody at 1:1000 dilution Lane 1: HepG2 whole cell lysates Lane 2: K562 whole cell lysates Lane 3: NCI-H460 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing A549 cells stained with AW5387(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AW5387, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OJ192088) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

### ALDH1A1 Antibody - Background

ALDH1A1 encodes a transcriptional regulator belonging to the SCY1-like family of kinase-like proteins. The protein has a divergent N-terminal kinase domain that is thought to be catalytically inactive, and can bind specific DNA sequences through its C-terminal domain. It activates transcription of the telomerase reverse transcriptase and DNA polymerase beta genes. The protein has been localized to the nucleus, and also to the cytoplasm and centrosomes during mitosis.

### ALDH1A1 Antibody - References

References for protein:

1. Gong, Y., et al. Oncogene 28(12):1549-1560(2009)
2. Burman, J.L., et al. J. Biol. Chem. 283(33):22774-22786(2008)
3. Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)

References for HepG2 cell line:

1. Knowles BB, et al. (1980). Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen. Science 209: 497-499.[ PubMed: 6248960].
2. Darlington GJ, et al. (1987). Growth and hepatospecific gene expression of human hepatoma cells in a defined medium. In Vitro Cell. Dev. Biol. 23: 349-354.[PubMed: 3034851].
3. Ihrke, G; Neufeld, EB; Meads, T; Shanks, MR; Cassio, D; Laurent, M; Schroer, TA; Pagano, RE et al. (1993). "WIF-B cells: an in vitro model for studies of hepatocyte polarity". Journal of Cell Biology 123 (6): 1761-1775. [PubMed:7506266].
4. Mersch-Sundermann, V.; Knasmüller, S.; Wu, X. J.; Darroudi, F.; Kassie, F. (2004). "Use of a human-derived liver cell line for the detection of cytoprotective, antigenotoxic and cogenotoxic agents". Toxicology 198 (1-3): 329-340. [PubMed:15138059].