

INSRR Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5413**Specification**

INSRR Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P14616
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=144 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

INSRR Antibody (Center) - Additional Information**Gene ID** 3645**Antigen Region**
668-702**Other Names**

Insulin receptor-related protein, IRR, IR-related receptor, Insulin receptor-related protein alpha chain, Insulin receptor-related protein beta chain, INSRR, IRR

Dilution

WB~~1:1000

IHC-P~~1:25

Target/Specificity

This INSRR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 668-702 amino acids from the Central region of human INSRR.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

INSRR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

INSRR Antibody (Center) - Protein Information

Name INSRR

Synonyms IRR

Function

Receptor with tyrosine-protein kinase activity. Functions as a pH sensing receptor which is activated by increased extracellular pH. Activates an intracellular signaling pathway that involves IRS1 and AKT1/PKB.

Cellular Location


Membrane; Single-pass type I membrane protein.

INSRR Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


INSRR Antibody (Center) - Images



All lanes : Anti-INSRR Antibody (Center) at 1:1000 dilution Lane 1: A549 whole cell lysates Lane 2: HeLa whole cell lysates Lane 3: MCF-7 whole cell lysates Lane 4: SH-SY5Y whole cell lysates Lane 5: SW480 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 144 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded H. kidney section using INSRR Antibody (Center)(Cat#AW5413). AW5413 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. breast carcinoma section using INSRR Antibody (Center)(Cat#AW5413). AW5413 was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

INSRR Antibody (Center) - Background

Receptor with tyrosine-protein kinase activity. Functions as a pH sensing receptor which is activated by increased extracellular pH. Activates an intracellular signaling pathway that involves IRS1 and AKT1/PKB.

INSRR Antibody (Center) - References

Haenze J.,et al.Horm. Metab. Res. 31:77-79(1999).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DBJ databases.
Shier P.,et al.J. Biol. Chem. 264:14605-14608(1989).
Greenman C.,et al.Nature 446:153-158(2007).