

MICA Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5477**Specification**

MICA Antibody (Center) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	Q29983
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=43,31 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

MICA Antibody (Center) - Additional Information**Gene ID** 100507436**Antigen Region**
68-97**Other Names**

MHC class I polypeptide-related sequence A, MIC-A, MICA {ECO:0000312|EMBL:CAI419071}

DilutionIF~~1:25
WB~~1:1000
IHC-P~~1:25
FC~~1:25**Target/Specificity**

This MICA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 68-97 amino acids from the Central region of human MICA.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MICA Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MICA Antibody (Center) - Protein Information

Name MICA {ECO:0000312|EMBL:CAI41907.1}

Function

Seems to have no role in antigen presentation. Acts as a stress-induced self-antigen that is recognized by gamma delta T-cells. Ligand for the KLRK1/NKG2D receptor. Binding to KLRK1 leads to cell lysis.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasm. Note=Expressed on the cell surface in gastric epithelium, endothelial cells and fibroblasts and in the cytoplasm in keratinocytes and monocytes. Infection with human adenovirus 5 suppresses cell surface expression due to the adenoviral E3-19K protein which causes retention in the endoplasmic reticulum

Tissue Location


Widely expressed with the exception of the central nervous system where it is absent. Expressed predominantly in gastric epithelium and also in monocytes, keratinocytes, endothelial cells, fibroblasts and in the outer layer of Hassal's corpuscles within the medulla of normal thymus. In skin, expressed mainly in the keratin layers, basal cells, ducts and follicles. Also expressed in many, but not all, epithelial tumors of lung, breast, kidney, ovary, prostate and colon. In thymomas, overexpressed in cortical and medullar epithelial cells. Tumors expressing MICA display increased levels of gamma delta T-cells.

MICA Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MICA Antibody (Center) - Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Hela (Human Cervical epithelial adenocarcinoma cell line) cells labeling MICA with AP8626c at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-rabbit IgG (NK179883) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on Hela cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DAPI (blue).




All lanes : Anti-MICA Antibody (Center) at 1:1000 dilution Lane 1: A431 whole cell lysates Lane 2: Daudi whole cell lysates Lane 3: Jurkat whole cell lysates Lane 4: MCF-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes : Anti-MICA Antibody (Center) at 1:2000 dilution Lane 1: Jurkat whole cell lysates Lane 2: Daudi whole cell lysates Lane 3: A431 whole cell lysates Lane 4: MCF-7 whole cell lysates Lane 5: U-87MG whole cell lysates Lane 6: THP-1 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Anti-MICA Antibody (Center) at 1:1000 dilution + HeLa whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Immunohistochemical analysis of paraffin-embedded H. hepatic carcinoma section using MICA Antibody (Center) (Cat#AP8626C). AP8626C was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. colorectal carcinoma section using MICA Antibody (Center)(Cat#AP8626C). AP8626C was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



AW5477 staining MICA in Human skin tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Flow cytometric analysis of SK-BR-3 cells using MICA Antibody (Center) (green, Cat#AP8626C) compared to an isotype control of rabbit IgG(blue). AP8626C was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

MICA Antibody (Center) - Background

MICA is the highly polymorphic MHC (HLA) class I chain-related gene A. The protein product is expressed on the cell surface, although unlike canonical class I molecules does not seem to associate with beta-2-microglobulin. It is thought that MICA functions as a stress-induced antigen that is broadly recognized by intestinal epithelial gamma delta T cells.

MICA Antibody (Center) - References

Bahram,S., et.al., Proc. Natl. Acad. Sci. U.S.A. 91 (14), 6259-6263 (1994) Klein,J.et.al., Proc. Natl. Acad. Sci. U.S.A. 91 (14), 6251-6252 (1994) Parham,P., et.al., J. Immunol. 142 (11), 3937-3950 (1989)