

SOX10 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AW5567

Specification

SOX10 Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Calculated MW Isotype Antigen Source

WB,E <u>P56693</u> <u>Q04888</u>, <u>A5A763</u>, <u>O55170</u> Human, Rat Mouse Rabbit Polyclonal H=50,31;M=50;R=50 KDa Rabbit IgG HUMAN

SOX10 Antibody (Center) - Additional Information

Gene ID 6663

Antigen Region 231-259

Other Names Transcription factor SOX-10, SOX10

Dilution WB~~1:2000

Target/Specificity This SOX10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 231-259 amino acids from the Central region of human SOX10.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions SOX10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SOX10 Antibody (Center) - Protein Information

Name SOX10

Function

Transcription factor that plays a central role in developing and mature glia (By similarity).



Specifically activates expression of myelin genes, during oligodendrocyte (OL) maturation, such as DUSP15 and MYRF, thereby playing a central role in oligodendrocyte maturation and CNS myelination (By similarity). Once induced, MYRF cooperates with SOX10 to implement the myelination program (By similarity). Transcriptional activator of MITF, acting synergistically with PAX3 (PubMed:>21965087). Transcriptional activator of MBP, via binding to the gene promoter (By similarity).

Cellular Location

Cytoplasm. Nucleus. Mitochondrion outer membrane {ECO:0000250|UniProtKB:Q04888}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q04888}; Cytoplasmic side {ECO:0000250|UniProtKB:Q04888}


Tissue Location Expressed in fetal brain and in adult brain, heart, small intestine and colon

SOX10 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- <u>Cell Culture</u>

SOX10 Antibody (Center) - Images



All lanes : Anti-SOX10 Antibody (Center) at 1:2000 dilution Lane 1: A375 whole cell lysate Lane 2: C6 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.