

PSMB7 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AW5615**Specification**

PSMB7 Antibody (Center) - Product Information

Application	WB, IHC,E
Primary Accession	Q99436
Other Accession	Q2TBP0 , P70195 , Q9JHW0
Reactivity	Human, Mouse
Predicted	Bovine, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	H=30;M=30;R=30 KDa
Isotype	Rabbit IgG
Antigen Source	HUMAN

PSMB7 Antibody (Center) - Additional Information**Gene ID** 5695**Antigen Region**
180-211**Other Names**

Proteasome subunit beta type-7, Macropain chain Z, Multicatalytic endopeptidase complex chain Z, Proteasome subunit Z, PSMB7, Z

DilutionWB~~1:8000
IHC~~1:25**Target/Specificity**

This PSMB7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 180-211 amino acids from the Central region of human PSMB7.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PSMB7 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PSMB7 Antibody (Center) - Protein Information**Name** PSMB7

Synonyms Z

Function

Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP- dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin- independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex). Within the 20S core complex, PSMB7 displays a trypsin-like activity.

Cellular Location

Cytoplasm. Nucleus. Note=Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9

Tissue Location


Expressed at a low level in colonic mucosa. Up- regulated in colorectal cancer tissues.

PSMB7 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PSMB7 Antibody (Center) - Images



All lanes : Anti-PSMB7 Antibody (Center) at 1:8000 dilution Lane 1: A375 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: MCF-7 whole cell lysate Lane 6: NIH/3T3 whole cell lysate Lane 7: Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 30 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AW5615 staining PSMB7 in human colorectal carcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

PSMB7 Antibody (Center) - Background

The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. This unit is responsible of the trypsin-like activity.

PSMB7 Antibody (Center) - References

Hisamatsu H.,et al.J. Exp. Med. 183:1807-1816(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Humphray S.J.,et al.Nature 429:369-374(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Lubec G.,et al.Submitted (DEC-2008) to UniProtKB.