

DANRE hoxc9a Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb10016a**Specification**

DANRE hoxc9a Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9YGS6
Reactivity	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	29782
Antigen Region	234-260

DANRE hoxc9a Antibody (C-term) - Additional Information**Other Names**

Homeobox protein Hox-C9a, Hox-C9, hoxc9a, hoxc9

Target/Specificity

This DANRE hoxc9a antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 234-260 amino acids from the C-terminal region of DANRE hoxc9a.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DANRE hoxc9a Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DANRE hoxc9a Antibody (C-term) - Protein Information

Name hoxc9a

Synonyms hoxc9

Function Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.


Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}.


DANRE hoxc9a Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DANRE hoxc9a Antibody (C-term) - Images

DANRE hoxc9a Antibody (C-term) (Cat. #Azbl0016a) western blot analysis in zebra fish brain tissue lysates (35ug/lane). This demonstrates the DANRE hoxc9a antibody detected the DANRE hoxc9a protein (arrow).



All lanes : Anti-DANRE hoxc9a Antibody (C-term) at 1:1000 dilution Lane 1: Zebrafish muscle lysate Lane 2: Zebrafish lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 30 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

DANRE hoxc9a Antibody (C-term) - Background

Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis (By similarity).

DANRE hoxc9a Antibody (C-term) - References

Amores A., et al. Science 282:1711-1714(1998).
Corredor-Adamez M., et al. Evol. Dev. 7:362-375(2005).
Prince V.E., et al. Development 125:407-420(1998).