

(DANRE) gapdh Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb18698c

Specification

(DANRE) gapdh Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q5XJ10
Other Accession	P04406
Reactivity	Human
Predicted	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	273-298

(DANRE) gapdh Antibody (C-term) - Additional Information

Gene ID 317743

Other Names

Glyceraldehyde-3-phosphate dehydrogenase, Peptidyl-cysteine S-nitrosylase GAPDH, 2699-, gapdh {ECO:0000312|ZFIN:ZDB-GENE-030115-1}

Target/Specificity

This (DANRE) gapdh antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 273-298 amino acids from the C-terminal region of DANRE gapdh.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(DANRE) gapdh Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) gapdh Antibody (C-term) - Protein Information

Name gapdh {ECO:0000312|ZFIN:ZDB-GENE-030115-1}

Function Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby

playing a role in glycolysis and nuclear functions, respectively. Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (By similarity). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

Cellular Location


Cytoplasm, cytosol {ECO:0000250|UniProtKB:P04797}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P04797}. Nucleus {ECO:0000250|UniProtKB:P04797}

(DANRE) gapdh Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(DANRE) gapdh Antibody (C-term) - Images



Western blot analysis of lysate from HUVEC cell line, using (DANRE) gapdh Antibody (C-term)(Cat. #Azb18698c). Azb18698c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

(DANRE) gapdh Antibody (C-term) - Background

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate. Modulates the

organization and assembly of the cytoskeleton. Also participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S- nitrosylation of nuclear target proteins (By similarity).