

(DANRE) edc3 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # Azb18719a

Specification

(DANRE) edc3 Antibody (Center) - Product Information

Application WB,E **Primary Accession** Q502M5 Reactivity Zebrafish Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 54852 Antigen Region 162-197

(DANRE) edc3 Antibody (Center) - Additional Information

Other Names

Enhancer of mRNA-decapping protein 3, LSM16 protein homolog, YjeF domain-containing protein 1, edc3, lsm16, yjdc

Target/Specificity

This (DANRE) edc3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 162-197 amino acids from the Central region of (DANRE) edc3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(DANRE) edc3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) edc3 Antibody (Center) - Protein Information

Name edc3

Synonyms Ism16, yjdc

Function Binds single-stranded RNA. Involved in the process of mRNA degradation and in the positive regulation of mRNA decapping (By similarity).



Tel: 858.875.1900 Fax: 858.875.1999

Cellular Location


Cytoplasm, P-body. Note=Processing bodies (PB).

(DANRE) edc3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

(DANRE) edc3 Antibody (Center) - Images



Western blot analysis of lysate from zebra fish muscle tissue lysate, using (DANRE) edc3 Antibody (Center)(Cat. #Azb18719a). Azb18719a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

(DANRE) edc3 Antibody (Center) - Background

Binds single-stranded RNA. In the process of mRNA degradation, may play a role in mRNA decapping (By similarity).

(DANRE) edc3 Antibody (Center) - References

Lemeer S., et al.J. Proteome Res. 7:1555-1564(2008).