10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

(DANRE) stk10 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # Azb19128c

Specification

(DANRE) stk10 Antibody (Center) - Product Information

Application	**WB,E**
Primary Accession	Q7SY52
Reactivity	**Zebrafish**
Host	**Rabbit**
Clonality	**polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**113687**
Antigen Region	**342-376**

(DANRE) stk10 Antibody (Center) - Additional Information

Gene ID 394108

Other Names
Serine/threonine-protein kinase 10, stk10

Target/Specificity
This DANRE stk10 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 342-376 amino acids from the Central region of DANRE stk10.

Dilution
WB~~1:2000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
(DANRE) stk10 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) stk10 Antibody (Center) - Protein Information

Name stk10

Function May act as a polo kinase kinase by mediating phosphorylation of plk1.

Cellular Location

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999


Cell membrane; Peripheral membrane protein

(DANRE) stk10 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

(DANRE) stk10 Antibody (Center) - Images



Anti-stk10 Antibody (Center)at 1:2000 dilution + ZF4 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 114 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

(DANRE) stk10 Antibody (Center) - Background

May act as a polo kinase kinase by mediating phosphorylation of plk1.