

# A4GALT Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP10116a

## **Specification**

# A4GALT Antibody (N-term) Blocking peptide - Product Information

Primary Accession Q9NPC4
Other Accession NP\_059132.1

# A4GALT Antibody (N-term) Blocking peptide - Additional Information

## Gene ID 53947

#### **Other Names**

Lactosylceramide 4-alpha-galactosyltransferase, Alpha-1, 4-N-acetylglucosaminyltransferase, Alpha-1, 4-galactosyltransferase, Alpha4Gal-T1, CD77 synthase, Globotriaosylceramide synthase, Gb3 synthase, P1/Pk synthase, UDP-galactose:beta-D-galactosyl-beta1-R 4-alpha-D-galactosyltransferase, A4GALT, A14GALT, A4GALT1

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## A4GALT Antibody (N-term) Blocking peptide - Protein Information

### Name A4GALT

Synonyms A14GALT, A4GALT1

## **Function**

Catalyzes the transfer of galactose from UDP-alpha-D- galactose to lactosylceramide/beta-D-galactosyl-(1->4)-beta-D-glucosyl- (1<->1)-ceramide(d18:1(4E)) to produce globotriaosylceramide/globoside Gb3Cer (d18:1(4E)) (PubMed:<a href="http://www.uniprot.org/citations/10748143" target="\_blank">10748143</a>). Also able to transfer galactose to galactosylceramide/beta-D-Gal-(1<->1')-Cer (PubMed:<a href="http://www.uniprot.org/citations/10748143" target="\_blank">10748143</a>). Globoside Gb3Cer is a glycosphingolipid of the globo serie, one of the major types of neutral root structures of glycosphingolipids, that constitute a significant portion of mammalian cell membranes (Probable). Globotriaosylceramide/globoside Gb3Cer in blood and tissue cell membranes is the antigen Pk of blood histogroup P (PubMed:<a href="http://www.uniprot.org/citations/10747952" target="\_blank">10747952</a>).



# **Cellular Location**

Golgi apparatus membrane; Single- pass type II membrane protein

#### **Tissue Location**

Ubiquitous. Highly expressed in kidney, heart, spleen, liver, testis and placenta

## A4GALT Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

A4GALT Antibody (N-term) Blocking peptide - Images

# A4GALT Antibody (N-term) Blocking peptide - Background

The protein encoded by this gene catalyzes the transfer ofgalactose to lactosylceramide to form globotriaosylceramide, whichhas been identified as the P(k) antigen of the P blood groupsystem. The encoded protein, which is a type II membrane proteinfound in the Golgi, is also required for the synthesis of thebacterial verotoxins receptor.

# A4GALT Antibody (N-term) Blocking peptide - References

Zumbrun, S.D., et al. Infect. Immun. 78(11):4488-4499(2010)Shin, I.S., et al. BMB Rep. 42(5):310-314(2009)Lund, N., et al. Blood 113(20):4980-4991(2009)Okuda, T., et al. Glycobiology 18(12):1028-1035(2008)Hellberg, A., et al. BMC Genet. 6, 49 (2005):