

# **COASY Antibody (Center) Blocking peptide**

Synthetic peptide Catalog # BP10371c

### **Specification**

# **COASY Antibody (Center) Blocking peptide - Product Information**

Primary Accession <u>Q13057</u>

Other Accession <u>NP\_001035995.1</u>, <u>NP\_001035997.2</u>,

NP\_079509.5, NP\_001035994.1,

NP 001035996.1

# **COASY Antibody (Center) Blocking peptide - Additional Information**

#### Gene ID 80347

#### **Other Names**

Bifunctional coenzyme A synthase, CoA synthase, NBP, POV-2, Phosphopantetheine adenylyltransferase, Dephospho-CoA pyrophosphorylase, Pantetheine-phosphate adenylyltransferase, PPAT, Dephospho-CoA kinase, DPCK, Dephosphocoenzyme A kinase, DPCOAK, COASY

### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **COASY Antibody (Center) Blocking peptide - Protein Information**

### Name COASY (HGNC:29932)

#### **Function**

Bifunctional enzyme that catalyzes the fourth and fifth sequential steps of CoA biosynthetic pathway. The fourth reaction is catalyzed by the phosphopantetheine adenylyltransferase, coded by the coaD domain; the fifth reaction is catalyzed by the dephospho-CoA kinase, coded by the coaE domain. May act as a point of CoA biosynthesis regulation.

## **Cellular Location**

Cytoplasm. Mitochondrion matrix. Note=The protein is mainly present in the mitochondrial matrix, probably anchored to the inner mitochondrial membrane, but is also present in cell lysate

### **Tissue Location**

Expressed in all tissues examined including brain, heart, skeletal muscle, colon, thymus, spleen, kidney, liver, small intestine, placenta, lung and peripheral blood leukocyte. Lowest expression in



peripheral blood leukocytes and highest in kidney and liver. Isoform 2 is expressed mainly in the brain

## **COASY Antibody (Center) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

### Blocking Peptides

## **COASY Antibody (Center) Blocking peptide - Images**

# COASY Antibody (Center) Blocking peptide - Background

Biosynthesis of coenzyme A (CoA) from pantothenic acid(vitamin B5) is an essential universal pathway in prokaryotes andeukaryotes. COASY is a bifunctional enzyme that catalyzes the 2last steps in CoA synthesis. These activities are performed by 2separate enzymes, phosphopantetheine adenylyltransferase (PPAT; EC2.7.7.3) and dephospho-CoA kinase (DPCK; EC 2.7.1.24), inprokaryotes (Daugherty et al., 2002 [PubMed 11923312]).[supplied byOMIM].

### COASY Antibody (Center) Blocking peptide - References

Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)Breus, O., et al. Biochem. Biophys. Res. Commun. 385(4):581-585(2009)Hamaguchi, M., et al. Jpn. J. Clin. Oncol. 38(11):734-742(2008)Nemazanyy, I., et al. Biochem. Biophys. Res. Commun. 341(4):995-1000(2006)Oh, J.H., et al. Mamm. Genome 16(12):942-954(2005)