

TPSD1 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP10500b**Specification**

TPSD1 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [O9BZJ3](#)
Other Accession [NP_036349.1](#)

TPSD1 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 23430

Other Names

Tryptase delta, Delta-tryptase, HmMCP-3-like tryptase III, Mast cell mMCP-7-like, Tryptase-3, TPSD1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TPSD1 Antibody (C-term) Blocking Peptide - Protein Information

Name TPSD1

Function

Tryptase is the major neutral protease present in mast cells and is secreted upon the coupled activation-degranulation response of this cell type.

Cellular Location

Secreted. Note=Released from the secretory granules upon mast cell activation.

Tissue Location

Expressed in colon, lung, heart and synovial tissue. May be specific to mast cells.

TPSD1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

TPSD1 Antibody (C-term) Blocking Peptide - Images**TPSD1 Antibody (C-term) Blocking Peptide - Background**

Tryptases comprise a family of trypsin-like serineproteases, the peptidase family S1. Tryptases are enzymatically active only as heparin-stabilized tetramers, and they are resistant to all known endogenous proteinase inhibitors. Several tryptase genes are clustered on chromosome 16p13.3. These genes are characterized by several distinct features. They have a highly conserved 3' UTR and contain tandem repeat sequences at the 5' flank and 3' UTR which are thought to play a role in regulation of the mRNA stability. Although this gene may be an exception, most of the tryptase genes have an intron immediately upstream of the initiator Met codon, which separates the site of transcription initiation from protein coding sequence. This feature is characteristic of tryptases but is unusual in other genes. Tryptases have been implicated as mediators in the pathogenesis of asthma and other allergic and inflammatory disorders. TPSD1 was once considered to be a pseudogene, although it is now believed to be a functional gene that encodes a protein.

TPSD1 Antibody (C-term) Blocking Peptide - References

Jackson, N.E., et al. J. Biol. Chem. 283(49):34178-34187(2008) Caughey, G.H. J. Allergy Clin. Immunol. 117(6):1411-1414(2006) Wang, H.W., et al. J. Immunol. 169(9):5145-5152(2002) Caughey, G.H. Mol. Immunol. 38 (16-18), 1353-1357 (2002) :Soto, D., et al. Clin. Exp. Allergy 32(7):1000-1006(2002)