

**PMAT(Slc29a4) Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP1087b****Specification**

---

**PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q8R139](#)**PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 243328**Other Names**

Equilibrative nucleoside transporter 4, Solute carrier family 29 member 4, Slc29a4, Ent4

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1087b](/product/products/AP1087b) was selected from the Slc29a4 region of human PMAT(Slc29a4). A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Protein Information****Name** Slc29a4 {ECO:0000312|MGI:MGI:2385330}**Function**

Electrogenic voltage-dependent transporter that mediates the transport of a variety of endogenous bioactive amines, cationic xenobiotics and drugs (PubMed: [16873718](http://www.uniprot.org/citations/16873718), PubMed: [23255610](http://www.uniprot.org/citations/23255610)). Utilizes the physiologic inside-negative membrane potential as a driving force to facilitate cellular uptake of organic cations (By similarity). Functions as a Na(+)- and Cl(-)-independent bidirectional transporter (By similarity). Substrate transport is pH-dependent and enhanced under acidic condition, which is most likely the result of allosteric changes in the transporter structure (PubMed: [16873718](http://www.uniprot.org/citations/16873718)). Implicated in monoamine neurotransmitters uptake such as serotonin, dopamine, adrenaline/epinephrine, noradrenaline/norepinephrine, histamine and tyramine, thereby supporting a role in homeostatic regulation of aminergic neurotransmission in the central nervous

system (PubMed:<a href="http://www.uniprot.org/citations/23255610" target="\_blank">23255610</a>). Also responsible for the uptake of bioactive amines and drugs through the blood-cerebrospinal fluid (CSF) barrier, from the CSF into choroid plexus epithelial cells, thereby playing a significant role in the clearance of cationic neurotoxins, xenobiotics and metabolic waste in the brain (PubMed:<a href="http://www.uniprot.org/citations/23255610" target="\_blank">23255610</a>). Involved in bidirectional transport of the purine nucleoside adenosine and plays a role in the regulation of extracellular adenosine concentrations in cardiac tissues, in particular during ischemia (PubMed:<a href="http://www.uniprot.org/citations/16873718" target="\_blank">16873718</a>). May be involved in organic cation uptake from the tubular lumen into renal tubular cells, thereby contributing to organic cation reabsorption in the kidney (PubMed:<a href="http://www.uniprot.org/citations/23255610" target="\_blank">23255610</a>). Also transports adenine and guanidine (PubMed:<a href="http://www.uniprot.org/citations/16873718" target="\_blank">16873718</a>).

#### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:Q7RTT9}; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Note=Localized to the apical blood-cerebrospinal fluid(CSF)-facing membrane of the choroid plexus epithelium

#### **Tissue Location**

Expressed in heart (PubMed:16873718). Expressed in choroid plexus (PubMed:23255610).

### **PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Images**

### **PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - Background**

PMAT(Slc29a4) is a member of the SLC29 family and encodes a plasma membrane protein with 11 transmembrane helices. This protein catalyzes the re-uptake of monoamines into presynaptic neurons, thus determining the intensity and duration of monoamine neural signaling. It has been shown to transport several compounds, including serotonin, dopamine, and the neurotoxin 1-methyl-4-phenylpyridinium.

### **PMAT(Slc29a4) Antibody (C-term) Blocking Peptide - References**

Barnes,K., Circ. Res. 99 (5), 510-519 (2006)Engel,K., J. Biol. Chem. 279 (48), 50042-50049 (2004)