

# **CEBPD Antibody (N-term) Blocking peptide**

Synthetic peptide Catalog # BP10950a

## **Specification**

## **CEBPD Antibody (N-term) Blocking peptide - Product Information**

Primary Accession

P49716

# CEBPD Antibody (N-term) Blocking peptide - Additional Information

**Gene ID 1052** 

#### **Other Names**

CCAAT/enhancer-binding protein delta, C/EBP delta, Nuclear factor NF-IL6-beta, NF-IL6-beta, CEBPD

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## CEBPD Antibody (N-term) Blocking peptide - Protein Information

### Name CEBPD

# **Function**

Transcription activator that recognizes two different DNA motifs: the CCAAT homology common to many promoters and the enhanced core homology common to many enhancers (PubMed:<a href="http://www.uniprot.org/citations/16397300" target="\_blank">16397300</a>). Important transcription factor regulating the expression of genes involved in immune and inflammatory responses (PubMed:<a href="http://www.uniprot.org/citations/1741402" target="\_blank">1741402</a>, PubMed:<a href="http://www.uniprot.org/citations/16397300" target="\_blank">16397300</a>). Transcriptional activator that enhances IL6 transcription alone and as heterodimer with CEBPB (PubMed:<a href="http://www.uniprot.org/citations/1741402" target="blank">1741402</a>).

### **Cellular Location**

Nucleus.

### **CEBPD Antibody (N-term) Blocking peptide - Protocols**



Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

### Blocking Peptides

# CEBPD Antibody (N-term) Blocking peptide - Images

# CEBPD Antibody (N-term) Blocking peptide - Background

The protein encoded by this intronless gene is a bZIPtranscription factor which can bind as a homodimer to certain DNAregulatory regions. It can also form heterodimers with the relatedprotein CEBP-alpha. The encoded protein is important in theregulation of genes involved in immune and inflammatory responses, and may be involved in the regulation of genes associated withactivation and/or differentiation of macrophages. The cytogeneticlocation of this locus has been reported as both 8p11 and 8q11.

# **CEBPD Antibody (N-term) Blocking peptide - References**

Zannetti, C., et al. J. Biol. Chem. 285(45):34773-34780(2010)Wang, J., et al. Proc. Natl. Acad. Sci. U.S.A. 107(37):16131-16136(2010)Hour, T.C., et al. Biochem. Pharmacol. 80(3):325-334(2010)Bennett, C.E., et al. Diab Vasc Dis Res 7(3):195-203(2010)Payne, V.A., et al. Biochem. J. 425(1):215-223(2010)