

**HDAC5 Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP1105b****Specification**

---

**HDAC5 Antibody (Center) Blocking Peptide - Product Information**

Primary Accession [O9UQL6](#)  
Other Accession [NP\\_005465](#)

**HDAC5 Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID** 10014

**Other Names**

Histone deacetylase 5, HD5, Antigen NY-CO-9, HDAC5, KIAA0600

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1105b](/product/products/AP1105b) was selected from the Center region of human HDAC5. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**HDAC5 Antibody (Center) Blocking Peptide - Protein Information**

**Name** HDAC5

**Synonyms** KIAA0600

**Function**

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Involved in muscle maturation by repressing transcription of myocyte enhancer MEF2C. During muscle differentiation, it shuttles into the cytoplasm, allowing the expression of myocyte enhancer factors. Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer. Serves as a corepressor of RARA and causes its deacetylation (PubMed:<http://www.uniprot.org/citations/28167758>). In

association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>).

**Cellular Location**

Nucleus. Cytoplasm. Note=Shuttles between the nucleus and the cytoplasm. In muscle cells, it shuttles into the cytoplasm during myocyte differentiation. The export to cytoplasm depends on the interaction with a 14-3-3 chaperone protein and is due to its phosphorylation at Ser-259 and Ser-498 by AMPK, CaMK1 and SIK1

**Tissue Location**

Ubiquitous.

**HDAC5 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**HDAC5 Antibody (Center) Blocking Peptide - Images****HDAC5 Antibody (Center) Blocking Peptide - Background**

Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the class II histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. It coimmunoprecipitates only with HDAC3 family member and might form multicomplex proteins. It also interacts with myocyte enhancer factor-2 (MEF2) proteins, resulting in repression of MEF2-dependent genes. This gene is thought to be associated with colon cancer. Two transcript variants encoding different isoforms have been noted for this gene.

**HDAC5 Antibody (Center) Blocking Peptide - References**

Berger, I., et al., J. Biol. Chem. 278(20):17625-17635 (2003).Zhang, C.L., et al., Mol. Cell. Biol. 22(20):7302-7312 (2002).Lemercier, C., et al., J. Biol. Chem. 277(24):22045-22052 (2002).Huang, Y., et al., Cancer Res. 62(10):2913-2922 (2002).Grozing, C.M., et al., Proc. Natl. Acad. Sci. U.S.A. 97(14):7835-7840 (2000).