

**GCDFP-15 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP11122b****Specification**

---

**GCDFP-15 Antibody (C-term) Blocking peptide - Product Information**Primary Accession [P12273](#)**GCDFP-15 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 5304**Other Names**

Prolactin-inducible protein, Gross cystic disease fluid protein 15, GCDFP-15, Prolactin-induced protein, Secretory actin-binding protein, SABP, gp17, PIP, GCDFP15, GPIP4

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**GCDFP-15 Antibody (C-term) Blocking peptide - Protein Information****Name** PIP**Synonyms** GCDFP15, GPIP4**Cellular Location**

Secreted.

**Tissue Location**

Expressed in pathological conditions of the mammary gland and in several exocrine tissues, such as the lacrimal, salivary, and sweat glands

**GCDFP-15 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**GCDFP-15 Antibody (C-term) Blocking peptide - Images**

**GCDFP-15 Antibody (C-term) Blocking peptide - Background**

Catalyzes the addition of fucose in alpha 1-6 linkage to the first GlcNAc residue, next to the peptide chains in N-glycans.

**GCDFP-15 Antibody (C-term) Blocking peptide - References**

Bryborn, M., et al. Allergy 65(2):220-228(2010)Wang, L.J., et al. Appl. Immunohistochem. Mol. Morphol. 17(6):505-511(2009)Yan, Z., et al. Diagn. Cytopathol. 37(7):475-478(2009)Debily, M.A., et al. PLoS ONE 4 (3), E4696 (2009) :Hassan, M.I., et al. J. Mol. Biol. 384(3):663-672(2008)