

# ZC3H8 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP11463a

### **Specification**

# ZC3H8 Antibody (N-term) Blocking peptide - Product Information

**Primary Accession** 

**Q8N5P1** 

### ZC3H8 Antibody (N-term) Blocking peptide - Additional Information

**Gene ID 84524** 

#### **Other Names**

Zinc finger CCCH domain-containing protein 8, ZC3H8, ZC3HDC8

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### ZC3H8 Antibody (N-term) Blocking peptide - Protein Information

Name ZC3H8

Synonyms ZC3HDC8

#### **Function**

Acts as a transcriptional repressor of the GATA3 promoter. Sequence-specific DNA-binding factor that binds to the 5'-AGGTCTC-3' sequence within the negative cis-acting element intronic regulatory region (IRR) of the GATA3 gene (By similarity). Component of the little elongation complex (LEC), a complex required to regulate small nuclear RNA (snRNA) gene transcription by RNA polymerase II and III (PubMed:<a href="http://www.uniprot.org/citations/23932780" target="\_blank">23932780</a>). Induces thymocyte apoptosis when overexpressed, which may indicate a role in regulation of thymocyte homeostasis.

#### **Cellular Location**

Nucleus. Note=Colocalizes with coilin in subnuclear cajal and histone locus bodies. Translocates in the LEC complex to cajal and histone locus bodies at snRNA genes in a ICE1-dependent manner. Associates to transcriptionally active chromatin at snRNA genes

# ZC3H8 Antibody (N-term) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

# ZC3H8 Antibody (N-term) Blocking peptide - Images

# ZC3H8 Antibody (N-term) Blocking peptide - Background

Acts as a transcriptional repressor of the GATA3 promoter. Sequence-specific DNA-binding factor that binds to the 5'-AGGTCTC-3' sequence within the negative cis-acting element intronic regulatory region (IRR) of the GATA3 gene (By similarity). Induces thymocyte apoptosis when overexpressed, which may indicate a role in regulation of thymocyte homeostasis.

# ZC3H8 Antibody (N-term) Blocking peptide - References

Lamesch, P., et al. Genomics 89(3):307-315(2007)Hwang, E.S., et al. Immunology 106(4):464-469(2002)Hwang, E.S., et al. J. Immunol. 169(1):248-253(2002)