

CACNG8 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP11619a

Specification

CACNG8 Antibody (N-term) Blocking peptide - Product Information

Primary Accession

Q8WXS5

CACNG8 Antibody (N-term) Blocking peptide - Additional Information

Gene ID 59283

Other Names

Voltage-dependent calcium channel gamma-8 subunit, Neuronal voltage-gated calcium channel gamma-8 subunit, Transmembrane AMPAR regulatory protein gamma-8, TARP gamma-8, CACNG8, CACNG6

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CACNG8 Antibody (N-term) Blocking peptide - Protein Information

Name CACNG8

Synonyms CACNG6

Function

Regulates the activity of L-type calcium channels that contain CACNA1C as pore-forming subunit (By similarity). Regulates the trafficking and gating properties of AMPA-selective glutamate receptors (AMPARs). Promotes their targeting to the cell membrane and synapses and modulates their gating properties by slowing their rates of activation, deactivation and desensitization and by mediating their resensitization. Does not show subunit-specific AMPA receptor regulation and regulates all AMPAR subunits.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q8VHW2}; Multi-pass membrane protein. Postsynaptic density membrane {ECO:0000250|UniProtKB:Q8VHW2}

Tissue Location

Detected in heart left ventricle.



CACNG8 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

CACNG8 Antibody (N-term) Blocking peptide - Images

CACNG8 Antibody (N-term) Blocking peptide - Background

The protein encoded by this gene is a type I transmembraneAMPA receptor regulatory protein (TARP). TARPs regulate bothtrafficking and channel gating of the AMPA receptors. This gene ispart of a functionally diverse eight-member protein subfamily of the PMP-22/EMP/MP20 family and is located in a cluster with twofamily members, a type II TARP and a calcium channel gamma subunit. The mRNA for this gene is believed to initiate translation from anon-AUG (CUG) start codon.

CACNG8 Antibody (N-term) Blocking peptide - References

Sager, C., et al. Neuroscience 158(1):45-54(2009)Correia, S.S., et al. Nat. Neurosci. 11(4):457-466(2008)Chen, R.S., et al. Cell Biochem. Biophys. 47(2):178-186(2007)Chu, P.J., et al. Gene 280 (1-2), 37-48 (2001) :Burgess, D.L., et al. Genomics 71(3):339-350(2001)