

MESDC2 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP11638b**Specification**

MESDC2 Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q14696](#)**MESDC2 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 23184**Other Names**

LDLR chaperone MESD, Mesoderm development candidate 2, Mesoderm development protein, Renal carcinoma antigen NY-REN-61, MESDC2, KIAA0081, MESD

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MESDC2 Antibody (C-term) Blocking peptide - Protein Information**Name** MESD ([HGNC:13520](#))**Synonyms** KIAA0081, MESDC2, MESDM**Function**

Chaperone specifically assisting the folding of beta- propeller/EGF modules within the family of low-density lipoprotein receptors (LDLRs) (PubMed:15014448). Acts as a modulator of the Wnt pathway through chaperoning the coreceptors of the canonical Wnt pathway, LRP5 and LRP6, to the plasma membrane (PubMed:17488095). Essential for specification of embryonic polarity and mesoderm induction. Plays an essential role in neuromuscular junction (NMJ) formation by promoting cell-surface expression of LRP4 (By similarity). May regulate phagocytosis of apoptotic retinal pigment epithelium (RPE) cells (By similarity).

Cellular Location

Endoplasmic reticulum Note=Released from apoptotic cells and shed photoreceptor outer segments. {ECO:0000250|UniProtKB:Q9ERE7}

MESDC2 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

MESDC2 Antibody (C-term) Blocking peptide - Images

MESDC2 Antibody (C-term) Blocking peptide - Background

Chaperone specifically assisting the folding of beta-propeller/EGF modules within the family of low-density lipoprotein receptors (LDLRs). Acts as a modulator of the Wnt pathway through chaperoning the coreceptors of the canonical Wnt pathway, LRP5 and LRP6, to the plasma membrane. Essential for specification of embryonic polarity and mesoderm induction.

MESDC2 Antibody (C-term) Blocking peptide - References

Murrills, R.J., et al. J. Cell. Biochem. 108(5):1066-1075(2009) Li, Y., et al. FEBS Lett. 580(22):5423-5428(2006) Veltman, I.M., et al. Hum. Mol. Genet. 14(14):1955-1963(2005) Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003) Hsieh, J.C., et al. Cell 112(3):355-367(2003)