

### Mash1 Antibody (L68) Blocking peptide

Synthetic peptide Catalog # BP11663a

### **Specification**

# Mash1 Antibody (L68) Blocking peptide - Product Information

**Primary Accession** 

P50553

### Mash1 Antibody (L68) Blocking peptide - Additional Information

Gene ID 429

#### **Other Names**

Achaete-scute homolog 1, ASH-1, hASH1, Class A basic helix-loop-helix protein 46, bHLHa46, ASCL1, ASH1, BHLHA46, HASH1

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### Mash1 Antibody (L68) Blocking peptide - Protein Information

Name ASCL1 (HGNC:738)

#### **Function**

Transcription factor that plays a key role in neuronal differentiation: acts as a pioneer transcription factor, accessing closed chromatin to allow other factors to bind and activate neural pathways. Directly binds the E box motif (5'-CANNTG-3') on promoters and promotes transcription of neuronal genes. The combination of three transcription factors, ASCL1, POU3F2/BRN2 and MYT1L, is sufficient to reprogram fibroblasts and other somatic cells into induced neuronal (iN) cells in vitro. Plays a role at early stages of development of specific neural lineages in most regions of the CNS, and of several lineages in the PNS. Essential for the generation of olfactory and autonomic neurons. Acts synergistically with FOXN4 to specify the identity of V2b neurons rather than V2a from bipotential p2 progenitors during spinal cord neurogenesis, probably through DLL4-NOTCH signaling activation. Involved in the regulation of neuroendocrine cell development in the glandular stomach (By similarity).

#### **Cellular Location**

Nucleus {ECO:0000250|UniProtKB:Q02067}.



Tel: 858.875.1900 Fax: 858.875.1999

# Mash1 Antibody (L68) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

Mash1 Antibody (L68) Blocking peptide - Images

### Mash1 Antibody (L68) Blocking peptide - Background

The protein encoded by this gene was identified by itsinteraction with the catalytic domain of protein kinase C-zeta. Theencoded protein contains an actin-binding site and an ATP-bindingsite. It is most closely related to twinfilin (PTK9), a conservedactin monomer-binding protein.

## Mash1 Antibody (L68) Blocking peptide - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)Han, S., et al. Hum. Immunol. 71(7):727-730(2010)Rajaraman, P., et al. Cancer Epidemiol. Biomarkers Prev. 19(5):1356-1361(2010)Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)Rajaraman, P., et al. Cancer Epidemiol. Biomarkers Prev. 18(5):1651-1658(2009)