

**COL8A2 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP11859b****Specification**

---

**COL8A2 Antibody (C-term) Blocking peptide - Product Information**Primary Accession [P25067](#)**COL8A2 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 1296**Other Names**

Collagen alpha-2(VIII) chain, Endothelial collagen, COL8A2

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**COL8A2 Antibody (C-term) Blocking peptide - Protein Information****Name** COL8A2**Function**

Macromolecular component of the subendothelium. Major component of the Descemet's membrane (basement membrane) of corneal endothelial cells. Also a component of the endothelia of blood vessels. Necessary for migration and proliferation of vascular smooth muscle cells and thus, has a potential role in the maintenance of vessel wall integrity and structure, in particular in atherogenesis (By similarity).

**Cellular Location**

Secreted, extracellular space, extracellular matrix, basement membrane

**Tissue Location**

Expressed primarily in the subendothelium of large blood vessels. Also expressed in arterioles and venules in muscle, heart, kidney, spleen, umbilical cord, liver and lung and is also found in connective tissue layers around hair follicles, around nerve bundles in muscle, in the dura of the optic nerve, in cornea and sclera, and in the perichondrium of cartilaginous tissues. In the kidney, expressed in mesangial cells, glomerular endothelial cells, and tubular epithelial cells. Also expressed in mast cells, and in astrocytes during the repair process. Expressed in Descemet's membrane

## **COL8A2 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **COL8A2 Antibody (C-term) Blocking peptide - Images**

## **COL8A2 Antibody (C-term) Blocking peptide - Background**

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes are involved in cellular defense against toxic, carcinogenic, and pharmacologically active electrophilic compounds. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase belonging to the alpha class genes that are located in a cluster mapped to chromosome 6. Genes of the alpha class are highly related and encode enzymes with glutathione peroxidase activity. However, during evolution, this alpha class gene diverged accumulating mutations in the active site that resulted in differences in substrate specificity and catalytic activity. The enzyme encoded by this gene catalyzes the double bond isomerization of precursors for progesterone and testosterone during the biosynthesis of steroid hormones. An additional transcript variant has been identified, but its full length sequence has not been determined.

## **COL8A2 Antibody (C-term) Blocking peptide - References**

Davila, S., et al. Genes Immun. 11(3):232-238(2010) Tars, K., et al. J. Mol. Biol. 397(1):332-340(2010) Moyer, A.M., et al. Cancer Epidemiol. Biomarkers Prev. 19(3):811-821(2010) Saito, A., et al. J. Hum. Genet. 54(6):317-323(2009) Starr, J.M., et al. Mech. Ageing Dev. 129(12):745-751(2008)