

# PCDHB6 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP12006a

# **Specification**

# PCDHB6 Antibody (N-term) Blocking peptide - Product Information

Primary Accession

**Q9Y5E3** 

# PCDHB6 Antibody (N-term) Blocking peptide - Additional Information

**Gene ID 56130** 

#### **Other Names**

Protocadherin beta-6, PCDH-beta-6, PCDHB6

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## PCDHB6 Antibody (N-term) Blocking peptide - Protein Information

Name PCDHB6 (HGNC:8691)

## **Function**

Calcium-dependent cell-adhesion protein involved in cells self-recognition and non-self discrimination. Thereby, it is involved in the establishment and maintenance of specific neuronal connections in the brain.

### **Cellular Location**

Cell membrane {ECO:0000250|UniProtKB:Q91XZ4}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q91XZ4}

# PCDHB6 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### Blocking Peptides

PCDHB6 Antibody (N-term) Blocking peptide - Images

PCDHB6 Antibody (N-term) Blocking peptide - Background







This gene is a member of the protocadherin beta genecluster, one of three related gene clusters tandemly linked onchromosome five. The gene clusters demonstrate an unusual genomicorganization similar to that of B-cell and T-cell receptor geneclusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmictail that deviates from others in the cadherin superfamily. Theextracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike the alpha and gammaclusters, the transcripts from these genes are made up of only onelarge exon, not sharing common 3' exons as expected. These neuralcadherin-like cell adhesion proteins are integral plasma membraneproteins. Their specific functions are unknown but they most likelyplay a critical role in the establishment and function of specificcell-cell neural connections.

# PCDHB6 Antibody (N-term) Blocking peptide - References

Frank, M., et al. Curr. Opin. Cell Biol. 14(5):557-562(2002)Vanhalst, K., et al. FEBS Lett. 495 (1-2), 120-125 (2001): Wu, Q., et al. Genome Res. 11(3):389-404(2001) Nollet, F., et al. J. Mol. Biol. 299(3):551-572(2000)Yagi, T., et al. Genes Dev. 14(10):1169-1180(2000)