

# PRMT8 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP1219b

# Specification

# PRMT8 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

<u>Q9NR22</u>

# PRMT8 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 56341

**Other Names** 

Protein arginine N-methyltransferase 8, 211-, Heterogeneous nuclear ribonucleoprotein methyltransferase-like protein 4, PRMT8, HRMT1L3, HRMT1L4

#### Target/Specificity

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP1219b>AP1219b</a> was selected from the C-term region of human PRMT8. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# PRMT8 Antibody (C-term) Blocking Peptide - Protein Information

Name PRMT8 (<u>HGNC:5188</u>)

# Synonyms HRMT1L3, HRMT1L4

Function

S-adenosyl-L-methionine-dependent and membrane-associated arginine methyltransferase that can both catalyze the formation of omega-N monomethylarginine (MMA) and asymmetrical dimethylarginine (aDMA) in proteins such as NIFK, myelin basic protein, histone H4, H2A and H2A/H2B dimer (PubMed:<a href="http://www.uniprot.org/citations/16051612" target="\_blank">16051612</a>, PubMed:<a href="http://www.uniprot.org/citations/16051612" target="\_blank">17925405</a>, PubMed:<a href="http://www.uniprot.org/citations/17925405" target="\_blank">26876602</a>, PubMed:<a href="http://www.uniprot.org/citations/26876602" target="\_blank">26876602</a>, PubMed:<a href="http://www.uniprot.org/citations/26879540" target="attp://www.uniprot.org/citations/26879540" target="attp://wwww.uniprot.org/citations/26879540"



href="http://www.uniprot.org/citations/18320585" target="\_blank">18320585</a>).

**Cellular Location** Cell membrane; Lipid-anchor; Cytoplasmic side

Tissue Location Brain-specific..

### PRMT8 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

#### PRMT8 Antibody (C-term) Blocking Peptide - Images

#### PRMT8 Antibody (C-term) Blocking Peptide - Background

Protein arginine methylation plays a role in signal transduction, RNA processing, transcriptional regulation, and DNA repair. PRMT8, a protein arginine N-methyltransferase most closely related to PRMT1, methylates the guanidino nitrogens of arginyl residues in some proteins. This protein associates with the plasma membrane following myristoylation and exhibits a brain-specific expression pattern, making it unique within this family of enzymes.

#### PRMT8 Antibody (C-term) Blocking Peptide - References

Lee, J., J. Biol. Chem. 280 (38), 32890-32896 (2005)