

HAX1 Antibody (C-term) Blocking peptide Synthetic peptide

Catalog # BP12311b

Specification

HAX1 Antibody (C-term) Blocking peptide - Product Information

Primary Accession

<u>000165</u>

HAX1 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 10456

Other Names HCLS1-associated protein X-1, HS1-associating protein X-1, HAX-1, HS1-binding protein 1, HSP1BP-1, HAX1, HS1BP1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

HAX1 Antibody (C-term) Blocking peptide - Protein Information

Name HAX1

Synonyms HS1BP1

Function

Recruits the Arp2/3 complex to the cell cortex and regulates reorganization of the cortical actin cytoskeleton via its interaction with KCNC3 and the Arp2/3 complex (PubMed:26997484). Slows down the rate of inactivation of KCNC3 channels (PubMed:26997484). Slows down the rate of inactivation of KCNC3 channels (PubMed:26997484). Slows down the rate of inactivation of KCNC3 channels (PubMed:26997484). Promotes

href="http://www.uniprot.org/citations/26997484" target="_blank">26997484). Promotes GNA13-mediated cell migration. Involved in the clathrin-mediated endocytosis pathway. May be involved in internalization of ABC transporters such as ABCB11. May inhibit CASP9 and CASP3. Promotes cell survival. May regulate intracellular calcium pools.

Cellular Location

Mitochondrion matrix. Endoplasmic reticulum Nucleus membrane. Cytoplasmic vesicle {ECO:0000250|UniProtKB:O35387}. Cytoplasm, cell cortex. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Sarcoplasmic reticulum {ECO:0000250|UniProtKB:Q7TSE9}. Cytoplasm, P-body [Isoform 3]: Cytoplasm. Nucleus Note=Predominantly cytoplasmic. Also detected in the nucleus when nuclear export is inhibited (in vitro). [Isoform 5]: Cytoplasm. Note=Predominantly



cytoplasmic

Tissue Location Ubiquitous. Up-regulated in oral cancers.

HAX1 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

HAX1 Antibody (C-term) Blocking peptide - Images

HAX1 Antibody (C-term) Blocking peptide - Background

The protein encoded by this gene is known to associatewith hematopoietic cell-specific Lyn substrate 1, a substrate ofSrc family tyrosine kinases. It also interacts with the product ofthe polycystic kidney disease 2 gene, mutations in which areassociated with autosomal-dominant polycystic kidney disease, andwith the F-actin-binding protein, cortactin. It was earlier thoughtthat this gene product is mainly localized in the mitochondria, however, recent studies indicate it to be localized in the cellbody. Mutations in this gene result in autosomal recessive severecongenital neutropenia, also known as Kostmann disease. Twotranscript variants encoding different isoforms have been found forthis gene.

HAX1 Antibody (C-term) Blocking peptide - References

Johns, H.L., et al. J. Gen. Virol. 91 (PT 11), 2677-2686 (2010) :Mekkawy, A.H., et al. Biochem. Biophys. Res. Commun. 399(4):738-743(2010)Han, J., et al. J. Biol. Chem. 285(29):22461-22472(2010)Germeshausen, M., et al. Haematologica 95(7):1207-1210(2010)Burnicka-Turek, O., et al. BMC Cell Biol. 11, 28 (2010) :