

POLR2G Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12458b

Specification

POLR2G Antibody (C-term) Blocking peptide - Product Information

Primary Accession [P62487](#)

POLR2G Antibody (C-term) Blocking peptide - Additional Information

Gene ID 5436

Other Names

DNA-directed RNA polymerase II subunit RPB7, RNA polymerase II subunit B7, DNA-directed RNA polymerase II subunit G, RNA polymerase II 19 kDa subunit, RPB19, POLR2G, RPB7

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

POLR2G Antibody (C-term) Blocking peptide - Protein Information

Name POLR2G

Synonyms RPB7

Function

DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Component of RNA polymerase II which synthesizes mRNA precursors and many functional non-coding RNAs. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB7 is part of a subcomplex with RPB4 that binds to a pocket formed by RPB1, RPB2 and RPB6 at the base of the clamp element. The RPB4-RPB7 subcomplex seems to lock the clamp via RPB7 in the closed conformation thus preventing double-stranded DNA to enter the active site cleft. The RPB4-RPB7 subcomplex binds single-stranded DNA and RNA (By similarity). Binds RNA.

Cellular Location

Nucleus.

POLR2G Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

POLR2G Antibody (C-term) Blocking peptide - Images**POLR2G Antibody (C-term) Blocking peptide - Background**

This gene encodes the seventh largest subunit of RNAPolymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. The protein functions in transcription initiation, and is also thought to help stabilize transcribing polymerase molecules during elongation.

POLR2G Antibody (C-term) Blocking peptide - References

Cojocaru, M., et al. Biochem. J. 409(1):139-147(2008)Ujvari, A., et al. Nat. Struct. Mol. Biol. 13(1):49-54(2006)Meka, H., et al. Nucleic Acids Res. 33(19):6435-6444(2005)Lehner, B., et al. Genomics 83(1):153-167(2004)Zhou, M., et al. Proc. Natl. Acad. Sci. U.S.A. 100(22):12666-12671(2003)