

PREX1 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12575b

Specification

PREX1 Antibody (C-term) Blocking peptide - Product Information

Primary Accession [Q8TCU6](#)

PREX1 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 57580

Other Names

Phosphatidylinositol 3, 5-trisphosphate-dependent Rac exchanger 1 protein, P-Rex1, PtdIns(3, 5)-dependent Rac exchanger 1, PREX1, KIAA1415

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PREX1 Antibody (C-term) Blocking peptide - Protein Information

Name PREX1

Synonyms KIAA1415

Function

Functions as a RAC guanine nucleotide exchange factor (GEF), which activates the Rac proteins by exchanging bound GDP for free GTP. Its activity is synergistically activated by phosphatidylinositol 3,4,5-trisphosphate and the beta gamma subunits of heterotrimeric G protein. May function downstream of heterotrimeric G proteins in neutrophils.

Cellular Location

Cytoplasm, cytosol. Cell membrane. Note=Mainly cytosolic. Some amount is apparently associated to the plasma membrane

Tissue Location

Mainly expressed in peripheral blood leukocytes and brain. Expressed at intermediate level in spleen and lymph nodes, and weakly expressed in other tissues

PREX1 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PREX1 Antibody (C-term) Blocking peptide - Images

PREX1 Antibody (C-term) Blocking peptide - Background

The protein encoded by this gene acts as a guaninenucleotide exchange factor for the RHO family of small GTP-binding proteins (RACs). It has been shown to bind to and activate RAC1 by exchanging bound GDP for free GTP. The encoded protein, which is found mainly in the cytoplasm, is activated by phosphatidylinositol-3,4,5-trisphosphate and the beta-gamma subunits of heterotrimeric G proteins.

PREX1 Antibody (C-term) Blocking peptide - References

Lewis, J.P., et al. Genomics 96(4):211-219(2010)
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010)
Nie, B., et al. Cell. Signal. 22(5):770-782(2010)
Carretero-Ortega, J., et al. Mol. Pharmacol. 77(3):435-442(2010)
Ledezma-Sanchez, B.A., et al. Biochem. Biophys. Res. Commun. 391(4):1647-1652(2010)