

**CBLC Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP1258a****Specification**

---

**CBLC Antibody (C-term) Blocking peptide - Product Information**Primary Accession [Q9ULV8](#)**CBLC Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 23624**Other Names**

E3 ubiquitin-protein ligase CBL-C, 632-, RING finger protein 57, SH3-binding protein CBL-3, SH3-binding protein CBL-C, Signal transduction protein CBL-C, CBLC, CBL3, RNF57

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP1258a](/product/products/AP1258a) was selected from the C-term region of human Cbl. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CBLC Antibody (C-term) Blocking peptide - Protein Information****Name** CBLC**Synonyms** CBL3, RNF57**Function**

Acts as an E3 ubiquitin-protein ligase, which accepts ubiquitin from specific E2 ubiquitin-conjugating enzymes, and then transfers it to substrates promoting their degradation by the proteasome. Functionally coupled with the E2 ubiquitin-protein ligases UB2D1, UB2D2 and UB2D3. Regulator of EGFR mediated signal transduction; upon EGF activation, ubiquitinates EGFR. Isoform 1, but not isoform 2, inhibits EGF stimulated MAPK1 activation. Promotes ubiquitination of SRC phosphorylated at 'Tyr-419'. In collaboration with CD2AP may act as regulatory checkpoint for Ret signaling by modulating the rate of RET degradation after ligand activation; CD2AP converts it from an inhibitor to a promoter of RET degradation; the function limits the potency of GDNF on neuronal survival.

**Tissue Location**

Ubiquitous..

**CBLC Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**CBLC Antibody (C-term) Blocking peptide - Images****CBLC Antibody (C-term) Blocking peptide - Background**

Cbl proteins are a family of ubiquitin protein ligases (E3s) that negatively regulate signaling by targeting activated tyrosine kinases for degradation. Cbl- c is the most recently cloned member of the Cbl proteins and is expressed only in epithelial cells (the other Cbl proteins are ubiquitously expressed). Cbl-c, like the other mammalian Cbl proteins, can ubiquitinate the activated EGFR and target it for degradation. Through interactions with proteins containing SRC homology-2 (SH2) and SH3 domains, CBL proteins modulate downstream cell signaling.

**CBLC Antibody (C-term) Blocking peptide - References**

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002).Keane, M.M., et al., Oncogene 18(22):3365-3375 (1999).Kim, M., et al., Gene 239(1):145-154 (1999).