

COPS7A Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12810b**Specification**

COPS7A Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [Q9UBW8](#)**COPS7A Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 50813**Other Names**

COP9 signalosome complex subunit 7a, SGN7a, Signalosome subunit 7a, Dermal papilla-derived protein 10, JAB1-containing signalosome subunit 7a, COPS7A, CSN7A, DERP10

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

COPS7A Antibody (C-term) Blocking peptide - Protein Information**Name** COPS7A**Synonyms** CSN7A, DERP10**Function**

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Widely expressed. Expressed at high level in brain, heart and skeletal muscle.

COPS7A Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

COPS7A Antibody (C-term) Blocking peptide - Images

COPS7A Antibody (C-term) Blocking peptide - Background

Component of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, JUN, I-kappa-B-alpha/NFKBIA, ITPK1 and IRF8/ICSBP, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

COPS7A Antibody (C-term) Blocking peptide - References

Matsuoka, S., et al. Science 316(5828):1160-1166(2007)Stelzl, U., et al. Cell 122(6):957-968(2005)Obuse, C., et al. Nat. Cell Biol. 6(11):1135-1141(2004)Wolf, D.A., et al. Nat. Cell Biol. 5(12):1029-1033(2003)Groisman, R., et al. Cell 113(3):357-367(2003)