

**VCC1 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP13104b****Specification**

---

**VCC1 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q6UXB2](#)**VCC1 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 284340**Other Names**

VEGF coregulated chemokine 1, C-X-C motif chemokine 17, Dendritic cell and monocyte chemokine-like protein, DMC, CXCL17, VCC1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13104b was selected from the C-term region of VCC1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**VCC1 Antibody (C-term) Blocking Peptide - Protein Information****Name** CXCL17**Synonyms** VCC1**Function**

Chemokine that acts as a chemoattractant for monocytes, macrophages and dendritic cells (PubMed: [16455961](http://www.uniprot.org/citations/16455961), PubMed: [23115081](http://www.uniprot.org/citations/23115081)). Plays a role in angiogenesis and possibly in the development of tumors (PubMed: [16989774](http://www.uniprot.org/citations/16989774), PubMed: [23115081](http://www.uniprot.org/citations/23115081)). Acts as an anti-inflammatory in the stomach (PubMed: [23115081](http://www.uniprot.org/citations/23115081)). May play a role in the innate defense against infections (PubMed: [17307946](http://www.uniprot.org/citations/17307946)). Activates the C-X-C chemokine receptor GPR35 to induce a rapid and transient rise in the level of

intracellular calcium ions (PubMed:<a href="http://www.uniprot.org/citations/25411203" target="\_blank">25411203</a>).

**Cellular Location**

Secreted.

**Tissue Location**

Detected in trachea, stomach, lung and skeletal muscle. Detected in intestine and in normal and asthmatic lung (at protein level). Breast tumors showed 3- to 24-fold up-regulation

**VCC1 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**VCC1 Antibody (C-term) Blocking Peptide - Images****VCC1 Antibody (C-term) Blocking Peptide - Background**

VCC1 plays a role in angiogenesis and possibly in the development of tumors. May be a housekeeping chemokine regulating recruitment of nonactivated blood monocytes and immature dendritic cells into tissues. May play a role in the innate defense against infections.

**VCC1 Antibody (C-term) Blocking Peptide - References**

Mu, X., et al. Acta Biochim. Biophys. Sin. (Shanghai) 41(8):631-637(2009)Weinstein, E.J., et al. Biochem. Biophys. Res. Commun. 350(1):74-81(2006)Pisabarro, M.T., et al. J. Immunol. 176(4):2069-2073(2006)Zlotnik, A., et al. Genome Biol. 7 (12), 243 (2006) :Zhang, Z., et al. Protein Sci. 13(10):2819-2824(2004)