

#### MSRB3 Antibody (N-term ) Blocking peptide Synthetic peptide

Catalog # BP13112a

# Specification

# MSRB3 Antibody (N-term ) Blocking peptide - Product Information

Primary Accession

<u>Q8IXL7</u>

# MSRB3 Antibody (N-term ) Blocking peptide - Additional Information

Gene ID 253827

Other Names Methionine-R-sulfoxide reductase B3, MsrB3, 184-, MSRB3

#### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13112a was selected from the N-term region of MSRB3. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage** Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# MSRB3 Antibody (N-term ) Blocking peptide - Protein Information

### Name MSRB3

Function

Catalyzes the reduction of free and protein-bound methionine sulfoxide to methionine. Isoform 2 is essential for hearing.

Cellular Location [Isoform 1]: Endoplasmic reticulum.

Tissue Location Widely expressed.

# MSRB3 Antibody (N-term ) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

#### Blocking Peptides

## MSRB3 Antibody (N-term ) Blocking peptide - Images

## MSRB3 Antibody (N-term ) Blocking peptide - Background

The protein encoded by this gene catalyzes the reduction of methionine sulfoxide to methionine. This enzyme acts as amonomer and requires zinc as a cofactor. Several transcriptvariants encoding two different isoforms have been found for thisgene. One of the isoforms localizes to mitochondria while the other localizes to endoplasmic reticula.

## MSRB3 Antibody (N-term ) Blocking peptide - References

Pillas, D., et al. PLoS Genet. 6 (2), E1000856 (2010) :Kwak, G.H., et al. BMB Rep 42(9):580-585(2009)Taungjaruwinai, W.M., et al. Am J Dermatopathol 31(5):427-431(2009)Lamesch, P., et al. Genomics 89(3):307-315(2007)Fortna, A., et al. PLoS Biol. 2 (7), E207 (2004) :