10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

PLAGL2 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP13129c

Specification

PLAGL2 Antibody (Center) Blocking Peptide - Product Information

Primary Accession Q9UPG8

PLAGL2 Antibody (Center) Blocking Peptide - Additional Information

Gene ID 5326

Other Names
Zinc finger protein PLAGL2, Pleiomorphic adenoma-like protein 2, PLAGL2, KIAA0198

Target/Specificity
The synthetic peptide sequence used to generate the antibody AP13129c was selected from the Center region of PLAGL2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format
Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions
This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PLAGL2 Antibody (Center) Blocking Peptide - Protein Information

Name PLAGL2

Synonyms KIAA0198

Function
Shows weak transcriptional activatory activity.

Cellular Location
Nucleus.

PLAGL2 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Blocking Peptides

PLAGL2 Antibody (Center) Blocking Peptide - Images

PLAGL2 Antibody (Center) Blocking Peptide - Background

Pleiomorphic adenoma gene-like 2 is a zinc-finger proteinthat recognizes DNA and/or RNA.

PLAGL2 Antibody (Center) Blocking Peptide - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Zheng, H., et al. Cancer Cell 17(5):497-509(2010)Wezensky, S.J., et al. Gene 452(1):22-34(2010)Yang, Y.S., et al. Am. J. Physiol. Lung Cell Mol. Physiol. 297 (3), L455-L466 (2009) :Guo, Y., et al. Biochem. Biophys. Res. Commun. 374(3):570-575(2008)