

PLS1 Antibody (C-term) Blocking Peptide Synthetic peptide Catalog # BP13147b

Specification

PLS1 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

<u>Q14651</u>

PLS1 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 5357

Other Names Plastin-1, Intestine-specific plastin, I-plastin, PLS1

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13147b was selected from the C-term region of PLS1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PLS1 Antibody (C-term) Blocking Peptide - Protein Information

Name PLS1

Function Actin-bundling protein. In the inner ear, it is required for stereocilia formation. Mediates liquid

packing of actin filaments that is necessary for stereocilia to grow to their proper dimensions.

Cellular Location Cytoplasm {ECO:0000250|UniProtKB:Q3V0K9}. Cell projection, stereocilium {ECO:0000250|UniProtKB:Q3V0K9}

Tissue Location

In small intestine, colon, and kidney; relatively lower levels of expression are detected in the lung and stomach



PLS1 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• **Blocking Peptides**

PLS1 Antibody (C-term) Blocking Peptide - Images

PLS1 Antibody (C-term) Blocking Peptide - Background

Plastins are a family of actin-binding proteins that areconserved throughout eukaryote evolution and expressed in mosttissues of higher eukaryotes. In humans, two ubiquitous plastinisoforms (L and T) have been identified. The protein encoded bythis gene is a third distinct plastin isoform, which isspecifically expressed at high levels in the small intestine. Alternatively spliced transcript variants varying in the 5' UTR, but encoding the same protein, have been found for this gene. Apseudogene of this gene is found on chromosome 11.

PLS1 Antibody (C-term) Blocking Peptide - References

Chafel, M.M., et al. Dev. Dyn. 203(2):141-151(1995)Lin, C.S., et al. Mol. Cell. Biol. 14(4):2457-2467(1994)Shibata, M., et al. J. Leukoc. Biol. 54(1):10-16(1993)Lin, C.S., et al. J. Biol. Chem. 268(4):2781-2792(1993)Zu, Y., et al. Biochemistry 29(4):1055-1062(1990)