

OR8I2 Antibody (C-term) Blocking peptide Synthetic peptide Catalog # BP13261b

## Specification

# **OR8I2 Antibody (C-term) Blocking peptide - Product Information**

Primary Accession

<u>Q8N0Y5</u>

# **OR8I2** Antibody (C-term) Blocking peptide - Additional Information

Gene ID 120586

Other Names Olfactory receptor 8l2, Olfactory receptor OR11-170, OR8l2

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13261b was selected from the C-term region of OR8I2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **OR8I2 Antibody (C-term) Blocking peptide - Protein Information**

Name OR8I2

Function Odorant receptor.

**Cellular Location** Cell membrane; Multi-pass membrane protein.

# **OR8I2 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

OR812 Antibody (C-term) Blocking peptide - Images



# OR8I2 Antibody (C-term) Blocking peptide - Background

Olfactory receptors interact with odorant molecules in thenose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a largefamily of G-protein-coupled receptors (GPCR) arising from singlecoding-exon genes. Olfactory receptors share a 7-transmembranedomain structure with many neurotransmitter and hormone receptorsand are responsible for the recognition and G protein-mediatedtransduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to theolfactory receptor genes and proteins for this organism isindependent of other organisms.

## OR8I2 Antibody (C-term) Blocking peptide - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)Fuchs, T., et al. Genomics 80(3):295-302(2002)