

# C10orf58 Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP13446b

### **Specification**

### C10orf58 Antibody (C-term) Blocking peptide - Product Information

**Primary Accession** 

Q9BRX8

## C10orf58 Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 84293** 

#### **Other Names**

Redox-regulatory protein FAM213A, Peroxiredoxin-like 2 activated in M-CSF stimulated monocytes, Protein PAMM, FAM213A, C10orf58, PAMM

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13446b was selected from the C-term region of C10orf58. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### C10orf58 Antibody (C-term) Blocking peptide - Protein Information

Name PRXL2A (HGNC:28651)

#### **Function**

Involved in redox regulation of the cell (PubMed:<a

href="http://www.uniprot.org/citations/26438880" target="\_blank">26438880</a>, PubMed:<a href="http://www.uniprot.org/citations/19951071" target="\_blank">19951071</a>). Acts as an antioxidant (PubMed:<a href="http://www.uniprot.org/citations/19951071" target="\_blank">19951071</a>). Acts as an antioxidant (PubMed:<a href="http://www.uniprot.org/citations/26438880" target="\_blank">19951071</a>). Inhibits TNFSF11-induced NFKB1 and JUN activation and osteoclast differentiation (PubMed:<a href="http://www.uniprot.org/citations/19951071" target="\_blank">19951071" target="\_blank">19951071</a>). May affect bone resorption and help to maintain bone mass (PubMed:<a href="http://www.uniprot.org/citations/19951071" target="\_blank">19951071</a>). Acts as a negative regulator of macrophage-mediated inflammation by inhibiting macrophage production of inflammatory cytokines, probably through suppression of the MAPK signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/26438880"



target="\_blank">26438880</a>).

#### **Cellular Location**

Cytoplasm. Secreted. Note=Secreted from mature adipocytes but not from preadipocytes.

#### **Tissue Location**

Expressed in CSF1 and TNFSF11-stimulated CD14(+) peripheral blood mononuclear cells (PBMCs)

## C10orf58 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

## • Blocking Peptides

C10orf58 Antibody (C-term) Blocking peptide - Images

C10orf58 Antibody (C-term) Blocking peptide - Background

The specific function of the protein remains unknown.

## C10orf58 Antibody (C-term) Blocking peptide - References

Castellano, S., et al. EMBO Rep. 5(1):71-77(2004)Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003)