

### LDHA Antibody (C-term) Blocking peptide Synthetic peptide

Catalog # BP13542b

## Specification

# LDHA Antibody (C-term) Blocking peptide - Product Information

Primary Accession

<u>P00338</u>

## LDHA Antibody (C-term) Blocking peptide - Additional Information

Gene ID 3939

#### **Other Names**

L-lactate dehydrogenase A chain, LDH-A, Cell proliferation-inducing gene 19 protein, LDH muscle subunit, LDH-M, Renal carcinoma antigen NY-REN-59, LDHA

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13542b was selected from the C-term region of LDHA. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### LDHA Antibody (C-term) Blocking peptide - Protein Information

Name LDHA (<u>HGNC:6535</u>)

#### Function

Interconverts simultaneously and stereospecifically pyruvate and lactate with concomitant interconversion of NADH and NAD(+).

Cellular Location Cytoplasm.

**Tissue Location** Predominantly expressed in anaerobic tissues such as skeletal muscle and liver.

### LDHA Antibody (C-term) Blocking peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

### <u>Blocking Peptides</u>

LDHA Antibody (C-term) Blocking peptide - Images

### LDHA Antibody (C-term) Blocking peptide - Background

The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of an aerobic glycolysis. The protein is found predominantly in muscletissue and belongs to the lactate dehydrogenase family. Mutations this gene have been linked to exertional myoglobinuria. Multipletranscript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene.

### LDHA Antibody (C-term) Blocking peptide - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Zhu, X., et al. Genet. Epidemiol. 34(2):171-187(2010)Zhuang, L., et al. Mod. Pathol. 23(1):45-53(2010)Zhao, Y.H., et al. Oncogene 28(42):3689-3701(2009)Koukourakis, M.I., et al. Oncology 77(5):285-292(2009)