

BTK Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP13546a**Specification**

BTK Antibody (N-term) Blocking peptide - Product InformationPrimary Accession [Q06187](#)**BTK Antibody (N-term) Blocking peptide - Additional Information****Gene ID** 695**Other Names**

Tyrosine-protein kinase BTK, Agammaglobulinemia tyrosine kinase, ATK, B-cell progenitor kinase, BPK, Bruton tyrosine kinase, BTK, AGMX1, ATK, BPK

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13546a was selected from the N-term region of BTK. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

BTK Antibody (N-term) Blocking peptide - Protein Information**Name** BTK**Synonyms** AGMX1, ATK, BPK**Function**

Non-receptor tyrosine kinase indispensable for B lymphocyte development, differentiation and signaling (PubMed:19290921). Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation (PubMed:19290921). After BCR engagement and activation at the plasma membrane, phosphorylates PLCG2 at several sites, igniting the downstream signaling pathway through calcium mobilization, followed by activation of the protein kinase C (PKC) family members (PubMed:11606584). PLCG2 phosphorylation is performed in close cooperation with the adapter protein B-cell linker protein

BLNK (PubMed:[11606584](http://www.uniprot.org/citations/11606584)). BTK acts as a platform to bring together a diverse array of signaling proteins and is implicated in cytokine receptor signaling pathways (PubMed:[16517732](http://www.uniprot.org/citations/16517732), PubMed:[17932028](http://www.uniprot.org/citations/17932028)). Plays an important role in the function of immune cells of innate as well as adaptive immunity, as a component of the Toll-like receptors (TLR) pathway (PubMed:[16517732](http://www.uniprot.org/citations/16517732)). The TLR pathway acts as a primary surveillance system for the detection of pathogens and are crucial to the activation of host defense (PubMed:[16517732](http://www.uniprot.org/citations/16517732)). Especially, is a critical molecule in regulating TLR9 activation in splenic B-cells (PubMed:[16517732](http://www.uniprot.org/citations/16517732), PubMed:[17932028](http://www.uniprot.org/citations/17932028)). Within the TLR pathway, induces tyrosine phosphorylation of TIRAP which leads to TIRAP degradation (PubMed:[16415872](http://www.uniprot.org/citations/16415872)). BTK also plays a critical role in transcription regulation (PubMed:[19290921](http://www.uniprot.org/citations/19290921)). Induces the activity of NF- κ B, which is involved in regulating the expression of hundreds of genes (PubMed:[19290921](http://www.uniprot.org/citations/19290921)). BTK is involved on the signaling pathway linking TLR8 and TLR9 to NF- κ B (PubMed:[19290921](http://www.uniprot.org/citations/19290921)). Acts as an activator of NLRP3 inflammasome assembly by mediating phosphorylation of NLRP3 (PubMed:[34554188](http://www.uniprot.org/citations/34554188)). Transiently phosphorylates transcription factor GTF2I on tyrosine residues in response to BCR (PubMed:[9012831](http://www.uniprot.org/citations/9012831)). GTF2I then translocates to the nucleus to bind regulatory enhancer elements to modulate gene expression (PubMed:[9012831](http://www.uniprot.org/citations/9012831)). ARID3A and NFAT are other transcriptional target of BTK (PubMed:[16738337](http://www.uniprot.org/citations/16738337)). BTK is required for the formation of functional ARID3A DNA-binding complexes (PubMed:[16738337](http://www.uniprot.org/citations/16738337)). There is however no evidence that BTK itself binds directly to DNA (PubMed:[16738337](http://www.uniprot.org/citations/16738337)). BTK has a dual role in the regulation of apoptosis (PubMed:[9751072](http://www.uniprot.org/citations/9751072)).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Nucleus Membrane raft {ECO:0000250|UniProtKB:P35991}. Note=In steady state, BTK is predominantly cytosolic. Following B-cell receptor (BCR) engagement by antigen, translocates to the plasma membrane through its PH domain Plasma membrane localization is a critical step in the activation of BTK. A fraction of BTK also shuttles between the nucleus and the cytoplasm, and nuclear export is mediated by the nuclear export receptor CRM1.

Tissue Location

Predominantly expressed in B-lymphocytes.

BTK Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

BTK Antibody (N-term) Blocking peptide - Images

BTK Antibody (N-term) Blocking peptide - Background

The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement.

BTK Antibody (N-term) Blocking peptide - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Ng, Y.Y., et al. Leukemia 24(9):1617-1630(2010) Segat, L., et al. Vaccine 28(10):2201-2206(2010) Marcotte, D.J., et al. Protein Sci. 19(3):429-439(2010) Liu, Z., et al. J. Immunol. 184(1):244-254(2010)