

ABHD4 Antibody (Center) Blocking peptide
Synthetic peptide
Catalog # BP13670c**Specification**

ABHD4 Antibody (Center) Blocking peptide - Product InformationPrimary Accession [Q8TB40](#)**ABHD4 Antibody (Center) Blocking peptide - Additional Information****Gene ID** 63874**Other Names**Abhydrolase domain-containing protein 4, 311-, Alpha/beta-hydrolase 4,
Lyso-N-acylphosphatidylethanolamine lipase, ABHD4**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13670c was selected from the Center region of ABHD4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ABHD4 Antibody (Center) Blocking peptide - Protein Information**Name** ABHD4 ([HGNC:20154](#))**Function**

Lysophospholipase selective for N-acyl phosphatidylethanolamine (NAPE). Contributes to the biosynthesis of N- acyl ethanolamines, including the endocannabinoid anandamide by hydrolyzing the sn-1 and sn-2 acyl chains from N-acyl phosphatidylethanolamine (NAPE) generating glycerophospho-N-acyl ethanolamine (GP-NAE), an intermediate for N-acyl ethanolamine biosynthesis. Hydrolyzes substrates bearing saturated, monounsaturated, polyunsaturated N-acyl chains. Shows no significant activity towards other lysophospholipids, including lysophosphatidylcholine, lysophosphatidylethanolamine and lysophosphatidylserine.

ABHD4 Antibody (Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ABHD4 Antibody (Center) Blocking peptide - Images

ABHD4 Antibody (Center) Blocking peptide - Background

Lysophospholipase selective for N-acyl phosphatidylethanolamine (NAPE). Contributes to the biosynthesis of N-acyl ethanolamines, including the endocannabinoid anandamide by hydrolyzing the sn-1 and sn-2 acyl chains from N-acyl phosphatidylethanolamine (NAPE) generating glycerophospho-N-acyl ethanolamine (GP-NAE), an intermediate for N-acyl ethanolamine biosynthesis. Hydrolyzes substrates bearing saturated, monounsaturated, polyunsaturated N-acyl chains. Shows no significant activity towards other lysophospholipids, including lysophosphatidylcholine, lysophosphatidylethanolamine and lysophosphatidylserine (By similarity).

ABHD4 Antibody (Center) Blocking peptide - References

Simon, G.M., et al. J. Biol. Chem. 281(36):26465-26472(2006)