

**STAT4 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP13722b****Specification**

---

**STAT4 Antibody (C-term) Blocking peptide - Product Information**Primary Accession [Q14765](#)**STAT4 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 6775**Other Names**

Signal transducer and activator of transcription 4, STAT4

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13722b was selected from the C-term region of STAT4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**STAT4 Antibody (C-term) Blocking peptide - Protein Information****Name** STAT4**Function**

Transcriptional regulator mainly expressed in hematopoietic cells that plays a critical role in cellular growth, differentiation and immune response (PubMed:<a href="http://www.uniprot.org/citations/8943379" target="\_blank">8943379</a>, PubMed:<a href="http://www.uniprot.org/citations/10961885" target="\_blank">10961885</a>, PubMed:<a href="http://www.uniprot.org/citations/37256972" target="\_blank">37256972</a>). Plays a key role in the differentiation of T-helper 1 cells and the production of interferon-gamma (PubMed:<a href="http://www.uniprot.org/citations/12213961" target="\_blank">12213961</a>, PubMed:<a href="http://www.uniprot.org/citations/35614130" target="\_blank">35614130</a>). Participates also in multiple neutrophil functions including chemotaxis and production of the neutrophil extracellular traps (By similarity). After IL12 binding to its receptor IL12RB2, STAT4 interacts with the intracellular domain of IL12RB2 and becomes tyrosine phosphorylated (PubMed:<a href="http://www.uniprot.org/citations/7638186" target="\_blank">7638186</a>, PubMed:<a href="http://www.uniprot.org/citations/10415122" target="\_blank">10415122</a>).

Phosphorylated STAT4 then homodimerizes and migrates to the nucleus where it can recognize STAT target sequences present in IL12 responsive genes. Although IL12 appears to be the predominant activating signal, STAT4 can also be phosphorylated and activated in response to IFN-gamma stimulation via JAK1 and TYK2 and in response to different interleukins including IL23, IL2 and IL35 (PubMed:<a href="http://www.uniprot.org/citations/11114383" target="\_blank">11114383</a>, PubMed:<a href="http://www.uniprot.org/citations/34508746" target="\_blank">34508746</a>). Transcription activation of IFN-gamma gene is mediated by interaction with JUN that forms a complex that efficiently interacts with the AP-1-related sequence of the IFN-gamma promoter (By similarity). In response to IFN- alpha/beta signaling, acts as a transcriptional repressor and suppresses IL5 and IL13 mRNA expression during response to T-cell receptor (TCR) activation (PubMed:<a href="http://www.uniprot.org/citations/26990433" target="\_blank">26990433</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Note=Translocated into the nucleus in response to phosphorylation.

#### **STAT4 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **STAT4 Antibody (C-term) Blocking peptide - Images**

#### **STAT4 Antibody (C-term) Blocking peptide - Background**

The protein encoded by this gene is a member of the STATfamily of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by their receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is essential for mediating responses to IL12 in lymphocytes, and regulating the differentiation of T helper cells.

#### **STAT4 Antibody (C-term) Blocking peptide - References**

Plant, D., et al. Ann. Rheum. Dis. 69(8):1548-1553(2010) Gestermann, N., et al. Genes Immun. 11(5):432-438(2010) Schuurhof, A., et al. Pediatr. Pulmonol. 45(6):608-613(2010) Vuong, M.T., et al. PLoS ONE 5 (5), E10559 (2010) :Glas, J., et al. PLoS ONE 5 (4), E10373 (2010) :