

# FCGR2C Antibody (C-term) Blocking peptide

Synthetic peptide Catalog # BP13730b

# **Specification**

# FCGR2C Antibody (C-term) Blocking peptide - Product Information

Primary Accession

# FCGR2C Antibody (C-term) Blocking peptide - Additional Information

**Gene ID 9103** 

#### **Other Names**

Low affinity immunoglobulin gamma Fc region receptor II-c, IgG Fc receptor II-c, CDw32, Fc-gamma RII-c, Fc-gamma-RIIc, FcRII-c, CD32, FCGR2C, CD32, FCG2, IGFR2

# **Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13730b was selected from the C-term region of FCGR2C. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

P31995

## **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# FCGR2C Antibody (C-term) Blocking peptide - Protein Information

Name FCGR2C

Synonyms CD32, FCG2, IGFR2

## **Function**

Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.

## **Cellular Location**

[Isoform IIC4]: Cytoplasm. [Isoform IIC2]: Cell membrane; Single-pass type I membrane protein

# **Tissue Location**

Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells



# FCGR2C Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

FCGR2C Antibody (C-term) Blocking peptide - Images

FCGR2C Antibody (C-term) Blocking peptide - Background

Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.