

MECP2 Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP13733a**Specification**

MECP2 Antibody (N-term) Blocking peptide - Product InformationPrimary Accession [P51608](#)**MECP2 Antibody (N-term) Blocking peptide - Additional Information****Gene ID** 4204**Other Names**

Methyl-CpG-binding protein 2, MeCp-2 protein, MeCp2, MECP2

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13733a was selected from the N-term region of MECP2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

MECP2 Antibody (N-term) Blocking peptide - Protein Information**Name** MECP2**Function**

Chromosomal protein that binds to methylated DNA. It can bind specifically to a single methyl-CpG pair. It is not influenced by sequences flanking the methyl-CpGs. Mediates transcriptional repression through interaction with histone deacetylase and the corepressor SIN3A. Binds both 5-methylcytosine (5mC) and 5-hydroxymethylcytosine (5hmC)- containing DNA, with a preference for 5-methylcytosine (5mC).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9Z2D6}. Note=Colocalized with methyl-CpG in the genome. Colocalized with TBL1X to the heterochromatin foci.

Tissue Location

Present in all adult somatic tissues tested.

MECP2 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

MECP2 Antibody (N-term) Blocking peptide - Images**MECP2 Antibody (N-term) Blocking peptide - Background**

DNA methylation is the major modification of eukaryotic genomes and plays an essential role in mammalian development. Human proteins MECP2, MBD1, MBD2, MBD3, and MBD4 comprise a family of nuclear proteins related by the presence in each of a methyl-CpG binding domain (MBD). Each of these proteins, with the exception of MBD3, is capable of binding specifically to methylated DNA. MECP2, MBD1 and MBD2 can also repress transcription from methylated gene promoters. In contrast to other MBD family members, MECP2 is X-linked and subject to X inactivation. MECP2 is dispensable in stem cells, but is essential for embryonic development. MECP2 gene mutations are the cause of most cases of Rett syndrome, a progressive neurologic developmental disorder and one of the most common causes of mental retardation in females. [provided by RefSeq].

MECP2 Antibody (N-term) Blocking peptide - References

Shapiro, J.R., et al. *Pediatr. Res.* 68(5):446-451(2010) Pintauro, M., et al. *Epilepsy Behav* (2010) In press : Jain, D., et al. *Pediatr. Neurol.* 43(1):35-40(2010) Harvey, C.G., et al. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 144B (3), 355-360 (2007) : Francke, U. *Nat Clin Pract Neurol* 2(4):212-221(2006)