

**MBL2 Antibody (C-term) Blocking peptide**  
**Synthetic peptide**  
**Catalog # BP13736b****Specification**

---

**MBL2 Antibody (C-term) Blocking peptide - Product Information**Primary Accession [P11226](#)**MBL2 Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 4153**Other Names**

Mannose-binding protein C, MBP-C, Collectin-1, MBP1, Mannan-binding protein, Mannose-binding lectin, MBL2, COLEC1, MBL

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody AP13736b was selected from the C-term region of MBL2. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**MBL2 Antibody (C-term) Blocking peptide - Protein Information****Name** MBL2 ([HGNC:6922](#))**Synonyms** COLEC1, MBL**Function**

Calcium-dependent lectin involved in innate immune defense (PubMed:<a href="http://www.uniprot.org/citations/35102342" target="\_blank">35102342</a>). Binds mannose, fucose and N-acetylglucosamine on different microorganisms and activates the lectin complement pathway. Binds to late apoptotic cells, as well as to apoptotic blebs and to necrotic cells, but not to early apoptotic cells, facilitating their uptake by macrophages. May bind DNA. Upon SARS coronavirus-2/SARS-CoV-2 infection, activates the complement lectin pathway which leads to the inhibition SARS-CoV-2 infection and a reduction of the induced inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/35102342" target="\_blank">35102342</a>).

**Cellular Location**

Secreted.

**Tissue Location**

Plasma protein produced mainly in the liver.

**MBL2 Antibody (C-term) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**MBL2 Antibody (C-term) Blocking peptide - Images****MBL2 Antibody (C-term) Blocking peptide - Background**

This gene encodes the soluble mannose-binding lectin or mannose-binding protein found in serum. The protein encoded belongs to the collectin family and is an important element in the innate immune system. The protein recognizes mannose and N-acetylglucosamine on many microorganisms, and is capable of activating the classical complement pathway. Deficiencies of this gene have been associated with susceptibility to autoimmune and infectious diseases.

**MBL2 Antibody (C-term) Blocking peptide - References**

Auriti, C., et al. Hum. Immunol. 71(11):1084-1088(2010) Fraser, D.A., et al. J. Immunol. 185(7):3932-3939(2010) de Wit, E., et al. Mamm. Genome (2010) In press : Filho, R.M., et al. Viral Immunol. 23(4):449-453(2010) Hu, Y., et al. Viral Immunol. 23(4):443-447(2010)