

# **DHRS7C Antibody (Center) Blocking peptide**

Synthetic peptide Catalog # BP13788c

## **Specification**

## **DHRS7C Antibody (Center) Blocking peptide - Product Information**

**Primary Accession** 

A6NNS2

## DHRS7C Antibody (Center) Blocking peptide - Additional Information

Gene ID 201140

#### **Other Names**

Dehydrogenase/reductase SDR family member 7C, 11--, DHRS7C

### Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13788c was selected from the Center region of DHRS7C. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **DHRS7C Antibody (Center) Blocking peptide - Protein Information**

Name DHRS7C (HGNC:32423)

#### **Function**

NADH-dependent oxidoreductase which catalyzes the oxidation of all-trans-retinol to all-trans-retinal. Plays a role in the regulation of cardiac and skeletal muscle metabolic functions. Maintains Ca(2+) intracellular homeostasis by repressing Ca(2+) release from the sarcoplasmic reticulum (SR) in myotubes, possibly through local alternations in NAD/NADH or retinol/retinal. Also plays a role in Ca(2+) homeostasis by controlling Ca(2+) overload in the cytosol and the SR in myotubes. Involved in glucose uptake into skeletal muscles and muscle performance by activating PI3K and mTORC2-mediated AKT1 phosphorylation signaling pathways, possibly through the action of its downstream catalytic product all-trans-retinoic acid.

## **Cellular Location**

Sarcoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q8CHS7}. Note=The N-terminus region encompasses a short hydrophobic sequence bound to the sarcoplasmic reticulum membrane, whereas the C-terminus catalytic domain faces the myoplasm In skeletal muscle,



enriched in the longitudinal sarcoplasmic reticulum. {ECO:0000250|UniProtKB:Q8CHS7}

# **DHRS7C Antibody (Center) Blocking peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

DHRS7C Antibody (Center) Blocking peptide - Images

**DHRS7C Antibody (Center) Blocking peptide - Background** 

DHRS7C is a putative oxidoreductase (Potential).