

# TNNC1 Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP13841a

# Specification

# TNNC1 Antibody (N-term) Blocking peptide - Product Information

Primary Accession

<u>P63316</u>

# TNNC1 Antibody (N-term) Blocking peptide - Additional Information

Gene ID 7134

**Other Names** Troponin C, slow skeletal and cardiac muscles, TN-C, TNNC1, TNNC

## Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13841a was selected from the N-term region of TNNC1. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **TNNC1** Antibody (N-term) Blocking peptide - Protein Information

Name TNNC1

Synonyms TNNC

Function

Troponin is the central regulatory protein of striated muscle contraction. Tn consists of three components: Tn-I which is the inhibitor of actomyosin ATPase, Tn-T which contains the binding site for tropomyosin and Tn-C. The binding of calcium to Tn-C abolishes the inhibitory action of Tn on actin filaments.

# TNNC1 Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

#### Blocking Peptides

## **TNNC1 Antibody (N-term) Blocking peptide - Images**

## TNNC1 Antibody (N-term) Blocking peptide - Background

Troponin is a central regulatory protein of striatedmuscle contraction, and together with tropomyosin, is located onthe actin filament. Troponin consists of 3 subunits: TnI, which isthe inhibitor of actomyosin ATPase; TnT, which contains the bindingsite for tropomyosin; and TnC, the protein encoded by this gene. The binding of calcium to TnC abolishes the inhibitory action ofTnI, thus allowing the interaction of actin with myosin, thehydrolysis of ATP, and the generation of tension. Mutations in thisgene are associated with cardiomyopathy dilated type 1Z. [providedby RefSeq].

### **TNNC1 Antibody (N-term) Blocking peptide - References**

Parvatiyar, M.S., et al. J. Biol. Chem. 285(36):27785-27797(2010)Swindle, N., et al. Biochemistry 49(23):4813-4820(2010)Dweck, D., et al. J. Biol. Chem. 285(23):17371-17379(2010)Bates, K.J., et al. Clin. Chem. 56(6):952-958(2010)Hershberger, R.E., et al. Circ Cardiovasc Genet 3(2):155-161(2010)