

Mouse Itk Antibody (N-term) Blocking peptide
Synthetic peptide
Catalog # BP13916a**Specification**

Mouse Itk Antibody (N-term) Blocking peptide - Product InformationPrimary Accession [Q03526](#)**Mouse Itk Antibody (N-term) Blocking peptide - Additional Information****Gene ID** 16428**Other Names**

Tyrosine-protein kinase ITK/TSK, IL-2-inducible T-cell kinase, Kinase EMT, Kinase TLK, T-cell-specific kinase, Itk, Emt, Tlk, Tsk

Target/Specificity

The synthetic peptide sequence used to generate the antibody AP13916a was selected from the N-term region of Mouse Itk. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mouse Itk Antibody (N-term) Blocking peptide - Protein Information**Name** Itk**Synonyms** Emt, Tlk, Tsk**Function**

Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylation lead to the recruitment of ITK to the cell membrane, in the vicinity of the stimulated TCR receptor, where it is phosphorylated by LCK. Phosphorylation leads to ITK autophosphorylation and full activation. Once activated, phosphorylates PLCG1, leading to the activation of this lipase and subsequent cleavage of its substrates. In turn, the endoplasmic reticulum releases calcium in the cytoplasm and the nuclear activator of activated T-cells (NFAT) translocates into the nucleus to perform its transcriptional duty. Phosphorylates 2 essential adapter proteins: the linker for activation of T-cells/LAT protein and LCP2. Then, a large number of

signaling molecules such as VAV1 are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Required for TCR- mediated calcium response in gamma-delta T-cells, may also be involved in the modulation of the transcriptomic signature in the Vgamma2- positive subset of immature gamma-delta T-cells (PubMed:23562159). Phosphorylates TBX21 at 'Tyr-525' and mediates its interaction with GATA3 (PubMed:15662016).

Cellular Location

Cytoplasm. Nucleus. Note=Localizes in the vicinity of cell surface receptors in the plasma membrane after receptor stimulation.

Tissue Location

Is detected in the thymus, lymph node and very faintly in the spleen, but is not detected in the liver, lung, kidney, heart, brain, intestine or testis. Expressed in T-lymphocytes and mast cells. It may also be expressed in natural killer cells

Mouse Itk Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Mouse Itk Antibody (N-term) Blocking peptide - Images**Mouse Itk Antibody (N-term) Blocking peptide - Background**

Itk may play a role in T-cell development, potentially in thymic selection.

Mouse Itk Antibody (N-term) Blocking peptide - References

Joseph, R.E., et al. J. Mol. Biol. 403(2):231-242(2010)Dierks, C., et al. Cancer Res. 70(15):6193-6204(2010)Grasis, J.A., et al. Mol. Cell. Biol. 30(14):3596-3609(2010)Xia, M., et al. J. Immunol. 184(12):6807-6814(2010)Sahu, N., et al. PLoS ONE 5 (6), E11348 (2010) :