

GTF2H1 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP14191a**Specification**

GTF2H1 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [P32780](#)**GTF2H1 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 2965**Other Names**

General transcription factor IIH subunit 1, Basic transcription factor 2 62 kDa subunit, BTF2 p62, General transcription factor IIH polypeptide 1, TFIIH basal transcription factor complex p62 subunit, GTF2H1, BTF2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

GTF2H1 Antibody (N-term) Blocking Peptide - Protein Information**Name** GTF2H1**Synonyms** BTF2**Function**

Component of the general transcription and DNA repair factor IIH (TFIIH) core complex, which is involved in general and transcription-coupled nucleotide excision repair (NER) of damaged DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. In NER, TFIIH acts by opening DNA around the lesion to allow the excision of the damaged oligonucleotide and its replacement by a new DNA fragment. In transcription, TFIIH has an essential role in transcription initiation. When the pre-initiation complex (PIC) has been established, TFIIH is required for promoter opening and promoter escape. Phosphorylation of the C-terminal tail (CTD) of the largest subunit of RNA polymerase II by the kinase module CAK controls the initiation of transcription.

Cellular Location

Nucleus.

GTF2H1 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

GTF2H1 Antibody (N-term) Blocking Peptide - Images**GTF2H1 Antibody (N-term) Blocking Peptide - Background**

Component of the core-TFIIF basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.

GTF2H1 Antibody (N-term) Blocking Peptide - References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Briggs, F.B., et al. Am. J. Epidemiol. 172(2):217-224(2010)Wu, W., et al. Lung Cancer 63(2):180-186(2009)Bethke, L., et al. J. Natl. Cancer Inst. 100(4):270-276(2008)Di Lello, P., et al. Biochemistry 44(21):7678-7686(2005)