

**POLR2I Antibody (N-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP14668a****Specification**

---

**POLR2I Antibody (N-term) Blocking Peptide - Product Information**

Primary Accession [P36954](#)

**POLR2I Antibody (N-term) Blocking Peptide - Additional Information**

**Gene ID** 5438

**Other Names**

DNA-directed RNA polymerase II subunit RPB9, RNA polymerase II subunit B9, DNA-directed RNA polymerase II subunit I, RNA polymerase II 145 kDa subunit, RPB145, POLR2I

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**POLR2I Antibody (N-term) Blocking Peptide - Protein Information**

**Name** POLR2I

**Function**

Core component of RNA polymerase II (Pol II), a DNA-dependent RNA polymerase which synthesizes mRNA precursors and many functional non-coding RNAs using the four ribonucleoside triphosphates as substrates. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. POLR2I/RPB9 is part of the upper jaw surrounding the central large cleft and thought to grab the incoming DNA template.

**Cellular Location**

Nucleus, nucleolus.

**POLR2I Antibody (N-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**POLR2I Antibody (N-term) Blocking Peptide - Images****POLR2I Antibody (N-term) Blocking Peptide - Background**

This gene encodes a subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. This subunit, in combination with two other polymerase subunits, forms the DNA binding domain of the polymerase, a groove in which the DNA template is transcribed into RNA. The product of this gene has two zinc finger motifs with conserved cysteines and the subunit does possess zinc binding activity. [provided by RefSeq].

**POLR2I Antibody (N-term) Blocking Peptide - References**

Matsuoka, S., et al. Science 316(5828):1160-1166(2007) Grimwood, J., et al. Nature 428(6982):529-535(2004) Zhou, M., et al. Proc. Natl. Acad. Sci. U.S.A. 100(22):12666-12671(2003) Kaehlcke, K., et al. Mol. Cell 12(1):167-176(2003) Hogan, T.H., et al. Biomed. Pharmacother. 57(1):41-48(2003)